

Spring 1987

INFLUENCE OF BIOGENIC AMINES ON BOVINE LUTEAL STEROIDOGENESIS

PAUL JEFFREY BATTISTA

University of New Hampshire, Durham

Follow this and additional works at: <https://scholars.unh.edu/dissertation>

Recommended Citation

BATTISTA, PAUL JEFFREY, "INFLUENCE OF BIOGENIC AMINES ON BOVINE LUTEAL STEROIDOGENESIS" (1987).
Doctoral Dissertations. 1502.
<https://scholars.unh.edu/dissertation/1502>

This Dissertation is brought to you for free and open access by the Student Scholarship at University of New Hampshire Scholars' Repository. It has been accepted for inclusion in Doctoral Dissertations by an authorized administrator of University of New Hampshire Scholars' Repository. For more information, please contact nicole.hentz@unh.edu.

INFORMATION TO USERS

While the most advanced technology has been used to photograph and reproduce this manuscript, the quality of the reproduction is heavily dependent upon the quality of the material submitted. For example:

- Manuscript pages may have indistinct print. In such cases, the best available copy has been filmed.
- Manuscripts may not always be complete. In such cases, a note will indicate that it is not possible to obtain missing pages.
- Copyrighted material may have been removed from the manuscript. In such cases, a note will indicate the deletion.

Oversize materials (e.g., maps, drawings, and charts) are photographed by sectioning the original, beginning at the upper left-hand corner and continuing from left to right in equal sections with small overlaps. Each oversize page is also filmed as one exposure and is available, for an additional charge, as a standard 35mm slide or as a 17"x 23" black and white photographic print.

Most photographs reproduce acceptably on positive microfilm or microfiche but lack the clarity on xerographic copies made from the microfilm. For an additional charge, 35mm slides of 6"x 9" black and white photographic prints are available for any photographs or illustrations that cannot be reproduced satisfactorily by xerography.

Order Number 8722426

Influence of biogenic amines on bovine luteal steroidogenesis

Battista, Paul Jeffrey, Ph.D.

University of New Hampshire, 1987

U·M·I
300 N. Zeeb Rd.
Ann Arbor, MI 48106

PLEASE NOTE:

In all cases this material has been filmed in the best possible way from the available copy. Problems encountered with this document have been identified here with a check mark ✓.

1. Glossy photographs or pages _____
2. Colored illustrations, paper or print _____
3. Photographs with dark background ✓ _____
4. Illustrations are poor copy _____
5. Pages with black marks, not original copy _____
6. Print shows through as there is text on both sides of page _____
7. Indistinct, broken or small print on several pages ✓ _____
8. Print exceeds margin requirements _____
9. Tightly bound copy with print lost in spine _____
10. Computer printout pages with indistinct print _____
11. Page(s) _____ lacking when material received, and not available from school or author.
12. Page(s) _____ seem to be missing in numbering only as text follows.
13. Two pages numbered _____. Text follows.
14. Curling and wrinkled pages _____
15. Dissertation contains pages with print at a slant, filmed as received _____
16. Other _____

University
Microfilms
International

INFLUENCE OF BIOGENIC AMINES ON
BOVINE LUTEAL STEROIDOGENESIS

BY

Paul Jeffrey Battista
A.A.S., State University of New York, Canton, 1979
B.S., West Virginia University, 1981
M.S., University of Maryland, 1983

DISSERTATION

Submitted to the University of New Hampshire
in Partial Fulfillment of
the Requirements for the Degree of

Doctor of Philosophy

in

Animal and Nutritional Sciences

May, 1987

This dissertation has been examined and approved.

William A. Condon

Dissertation director, William A. Condon
Chairperson, Department of Animal
and Nutritional Sciences

William E. Berndtson

William E. Berndtson, Associate
Professor of Animal and Nutritional
Sciences

Thomas Foxall

Thomas Foxall, Assistant
Professor of Animal and
Nutritional Sciences

Stacia A. Sower

Stacia A. Sower, Assistant
Professor of Zoology

Winsor H. Watson III

Winsor H. Watson III, Associate
Professor of Zoology

3/23/87
Date

TABLE OF CONTENTS

LIST OF TABLES.....	v
LIST OF FIGURES.....	vi
DRUG DESCRIPTION.....	vii
ABSTRACT.....	viii
SECTION	PAGE
I. REVIEW OF THE LITERATURE.....	1
Neural Control of Reproductive Function.....	1
Neural Innervation of the Mammalian Reproductive Tract and Ovary.....	2
II. THE CORPUS LUTEUM.....	4
III. LUTEOTROPISM.....	8
IV. FACTORS REGULATING OVARIAN STEROIDOGENESIS.....	9
Catecholamines.....	9
Serotonin and Melatonin.....	11
Histamine and Dopamine.....	12
Acetylcholine.....	12
Gamma-Aminobutyric Acid.....	13
Vasoactive Intestinal Peptide.....	13
Substance P.....	14
Opiates.....	14
Oxytocin and Vasopressin.....	15
Relaxin.....	16
Insulin and Insulin-Like Growth Factors.....	17
Epidermal Growth Factor.....	18
Gonadotropin Releasing-Hormone.....	18
Adenosine.....	19

Prostaglandins.....	20
Glucocorticoids.....	21
Catecholestrogens.....	22
V. MECHANISMS CONTROLLING STEROIDOGENESIS.....	23
Hormone-Receptor Interaction: Activation of Adenylate Cyclase.....	23
cAMP and cAMP-Dependent Protein Kinase.....	25
Regulation of Steroidogenesis: Protein Kinase A.....	25
Regulation of Steroidogenesis: Protein Kinase C.....	28
Progesterone Secretion.....	29
VI. REFERENCES.....	31
VII. EXPERIMENT ONE.....	61
Serotonin-Induced Stimulation of Progesterone Production by Cow Luteal Cells <u>In Vitro</u>	
VIII. EXPERIMENT TWO.....	83
Mechanisms Involved in the Action of Serotonin-Induced Stimulation of Progesterone Production by Bovine Luteal Cells <u>In Vitro</u>	
IX. EXPERIMENT THREE.....	105
A Role for Alternative Pathway Catecholamines in the Stimulation of Steroidogenesis in Cow Luteal Cells.	
X. EXPERIMENT FOUR.....	122
Biogenic Amine Stimulation of Bovine Luteal Progesterone Production <u>In Vivo</u>	
XI. SUMMARY AND SIGNIFICANCE OF RESULTS.....	141

LIST OF TABLES

	PAGE
EXPERIMENT ONE	
1. Effect of biogenic amines on progesterone production by dissociated luteal cells in a 2-h incubation.....	77
2. Effect of biogenic amines on LH-stimulated progesterone production by dissociated luteal cells in a 2-h incubation.....	78
EXPERIMENT TWO	
1. Effect of serotonin metabolites on progesterone production by dissociated luteal cells in short-term incubation.....	99
EXPERIMENT THREE	
1. Effect of α and β -adrenergic antagonists on primary and alternative pathway catecholamine-induced progesterone production.....	117
EXPERIMENT FOUR	
1. Effect of biogenic amines on mean plasma progesterone concentrations (ng/ml) over a 72-h treatment period.....	137

LIST OF FIGURES

EXPERIMENT ONE	PAGE
1. Effect of serotonin on progesterone production by dissociated luteal cells.....	80
2. Effect of Isoproterenol on luteal cell progesterone production.....	81
3. Effect of β -adrenergic antagonists and mianserin on Isoproterenol-stimulated progesterone production.....	82
 EXPERIMENT TWO	
1. Effect of 5-HT antagonists on 5-HT-induced progesterone production.....	101
2. Effect of the 5-HT ₁ agonist DPAT on luteal cell progesterone production.....	102
3. Effect of IBMX on basal progesterone production and progesterone production induced by 5-HT and DPAT either alone or in combination with 5-HT antagonists.....	103
4. 5-HT concentration of bovine luteal tissue as affected by the stage of the estrous cycle.....	104
 EXPERIMENT THREE	
1. Primary and alternative pathways for the formation of catecholamines.....	119
2. Effect of alternative pathway catecholamines on progesterone production by dissociated bovine luteal cells.....	120
3. Effect of noradrenaline and adrenaline on progesterone production by dissociated luteal cells.....	121
 EXPERIMENT FOUR	
1a. Time-course effect of noradrenaline and adrenaline on the <u>in vivo</u> stimulation of bovine luteal progesterone production.....	139
1b. Time-course effect of 5-HT on the <u>in vivo</u> stimulation of bovine luteal progesterone production.....	140

DRUG DESCRIPTION

2-Amino-6,7-dihydroxy-1,2,3,4-tetranaphthalene hydrobromide	nonspecific dopamine receptor agonist
Butoxamine	β_2 -adrenergic receptor antagonist
8-Hydroxydiethylpropylamine	specific serotonin-1 receptor agonist
3-Isobutyl-methylxanthine (IBMX)	phosphodiesterase inhibitor
Isoproterenol	nonspecific β -adrenergic receptor agonist
Ketanserin	specific serotonin-2 receptor antagonist
Mianserin	nonspecific serotonin receptor antagonist
Phentolamine	nonspecific α -adrenergic receptor antagonist
Practolol	β_1 -adrenergic receptor antagonist
Propranolol	nonspecific β -adrenergic receptor antagonist
Salbutamol	β_2 -adrenergic receptor agonist
Terbutaline	β_2 -adrenergic receptor agonist

ABSTRACT

Influence of Biogenic Amines on Bovine
Luteal Steroidogenesis

by

Paul Jeffrey Battista

University of New Hampshire, May 1987

To investigate the role of catecholamines and biogenic amines in the control of bovine luteal steroidogenesis, studies were conducted utilizing collagenase-dispersed luteal cells in a short-term incubation system. Studies designed to assess the role of catecholamines showed that the stimulatory response to the primary catecholamines norepinephrine and epinephrine was mediated by both β_1 - and β_2 -adrenergic receptors. The alternative pathway catecholamines octopamine, synephrine and deoxyepinephrine enhanced basal progesterone production, and this response was mediated by β -adrenergic receptors. Analysis of luteal tissue using high performance liquid chromatography showed nondetectable levels of norepinephrine and epinephrine. The in vivo administration of norepinephrine and epinephrine significantly stimulated luteal steroidogenesis without altering circulating levels of luteinizing hormone.

The addition of serotonin or 5-methoxytryptamine to luteal cell incubations increased the production of progesterone in a dose-dependent manner. The response to serotonin was mediated by a serotonin-1 receptor, which may not be associated with adenylate cyclase. Analysis of bovine luteal tissue using high performance liquid chromatography demonstrated the presence of serotonin, and showed that the concentration of serotonin varied with the stage of luteal development. Administration of serotonin in vivo resulted in significantly elevated levels of plasma progesterone, which were independent of changes in luteinizing hormone.

These results support the concept that a variety of biogenic amines influence bovine luteal steroidogenesis.

REVIEW OF THE LITERATURE

Neural Control of Reproductive Function

Most, if not all, reproductive functions are controlled by hormonal and neural influences. While the hormonal control of reproductive function has received the most attention (reviewed by Hansel et al., 1973; Niswender et al., 1985) the nervous system has been implicated in controlling a variety of reproductive processes including: gonadotropin secretion (reviewed by Naumenko, 1985), cyclicity (Hill, 1949, 1962; Gelderd and Pepper, 1979), prepubertal ovarian development (Aguado et al., 1982; Aguado and Ojeda, 1984a; Ben-Johnathan et al., 1984), follicular development (Brink and Grob, 1972; Grob, 1974; Burden, 1978; Curry et al., 1983), ovulation (reviewed by Bahr et al., 1974), ovarian steroidogenesis (Condon and Black, 1976; Godkin et al., 1977; Kawakami et al., 1981; Weiss et al., 1982), compensatory ovarian hypertrophy (Gerendai et al., 1978; Moger and Anakwe, 1986), pregnancy (reviewed by Burden, 1978), parturition (reviewed by Burden, 1978; Maltier, 1984; Legrand and Maltier, 1986), uterine and oviductal motility (reviewed by Black, 1979), uterine and ovarian blood flow (reviewed by Ford, 1985), puberty (reviewed by Foster et al., 1985), and seasonal reproduction (reviewed by Lincoln, 1984).

The following review of the literature is concerned with the relationship between the peripheral nervous system and the regulation of ovarian function. More specifically, the major focus of the following

literature review and the subsequent studies performed, is concerned with elucidating the role of biogenic amines in the control of steroidogenesis by the corpus luteum.

Neural Innervation of the Mammalian Reproductive Tract and Ovary

The mammalian reproductive tract is innervated by both sympathetic and parasympathetic components of the autonomic nervous system (reviewed by Bell, 1972; Black, 1974, 1979). Both noradrenergic and cholinergic nerve fibers innervate the vascular and nonvascular smooth muscle of the oviducts, uterus, cervix and vagina (reviewed by Papka et al., 1985). The sympathetic component of the autonomic nervous system innervating the reproductive tract arises from the hypogastric nerves (Brudin, 1965; Sjöberg, 1967; Marshall, 1970; Bell, 1972) while parasympathetic innervation arises from the pelvic nerve (Schofield, 1952; Carlson and DeFeo, 1965; Marshall, 1970). The mammalian oviduct also receives sympathetic innervation via the inferior, middle and superior ovarian nerves (Mitchell, 1938; Marshall, 1970; Bell, 1972). The reproductive system is more highly innervated by noradrenergic than cholinergic nerve fibers (Owman, 1967; Sjöberg, 1967; Black, 1974, 1979). The distribution of noradrenergic and cholinergic nerve fibers supplying the reproductive tract follows a similar pattern across mammalian species, although the degree of innervation varies widely among species (reviewed by Black, 1974, 1979; Papka et al., 1985).

As in the reproductive tract, the ovary is innervated by both noradrenergic and cholinergic nerve fibers (reviewed by Bahr et al., 1974). The ovary receives sympathetic innervation from two main sources, the superior ovarian nerve, which travels along the suspensory ligament and the plexus nerve which accompanies the ovarian artery in

the ovarian plexus (Lawrence and Burden, 1980). Nerve fibers of the ovarian plexus terminate primarily on blood vessels while those of the superior ovarian nerve terminate on blood vessels and the steroid producing cells of the interstitial gland (Lawrence and Burden, 1980). Sympathetic nerve fibers arising from the superior ovarian nerve innervate the theca but not the granulosa cell layer of ovarian follicles (Burden, 1972; Unsicker, 1974; Wallis, 1975, 1978; Stefenson et al., 1981). In addition, the corpus luteum is devoid of noradrenergic innervation (Burden, 1972; Unsicker, 1974; Black, 1979).

Cholinergic innervation of the mammalian ovary appears to arise from the vagus nerve (Hill, 1962; LePere et al., 1966; Burden and Lawrence, 1978). but parasympathetic innervation from the hypogastric (Stohr, 1957) and ovarian plexus (Burden et al., 1978) has also been demonstrated. Cholinergic innervation of the ovary is less developed but has the same distribution as noradrenergic fibers innervating blood vessels, ovarian stroma and the follicular wall, but not the corpus luteum (Jacobowitz and Wallach, 1967; Owman et al., 1975; Wallis et al., 1976; Burden and Lawrence, 1978; Black, 1979; Stefenson et al., 1981). As in the reproductive tract, the distribution of sympathetic and parasympathetic nerve fibers supplying the ovary tends to follow a similar pattern among species, although the degree of innervation does vary (Bahr et al., 1974; Burden, 1979; Stefenson et al., 1981).

Recently a variety of other biogenic amines and neuroactive peptides have been observed in mammalian reproductive organs. These compounds include dopamine (Farrar et al., 1980; Van Orden et al., 1980; Bahr and Ben-Johnathan 1981; Ben-Johnathan et al., 1984), epinephrine (Wurtman et al., 1963; Van Orden et al., 1980; Bahr and Ben-Johnathan 1985).

serotonin (Clausell and Soliman, 1978; Battista et al., 1987a), substance P (Makris et al., 1982; Dees et al., 1985; Ojeda et al., 1985; Papka et al., 1985), vasoactive intestinal peptide (Larsson et al., 1977; Alm et al., 1980a, b; Hakanson et al., 1982), gamma-aminobutyric acid (Del Rio and Caballero, 1980; Del Rio, 1981; Erdo et al., 1982; Schaffer and Hsueh, 1982), β -endorphin opiates (Tsong et al., 1982a,b; Alm et al., 1983; Lim et al., 1983; Lolait et al., 1985), neuropeptide tyrosine (Sternquist et al., 1983; Taurig et al., 1984; Papka et al., 1985), neuropeptide Y (Sternquist et al., 1983; Blank et al., 1986), oxytocin, oxytocin-associated neurophysin, vasopressin (Wathes and Swann, 1982; Wathes et al., 1983; Wathes, 1983), and relaxin (reviewed by Sherwood and Downing, 1983).

The role of these compounds, as well as norepinephrine and acetylcholine, in the control of ovarian steroidogenesis will be addressed in a subsequent portion of this literature review.

THE CORPUS LUTEUM

The mammalian corpus luteum (CL) is a transient endocrine gland which was first observed by Volcherus Colter in 1573 and later named as such by Marcello Malpighi in 1697 due to its yellow appearance in the cow. The early studies of Fraenkel and Cohn (1901) established that the CL was essential for maintenance of pregnancy in rabbits and that it appeared to meet the criteria of being an endocrine gland. In 1917 Loeb reported that guinea pigs required a CL for ova to induce development of the placenta. Later, Corner and Allen (1929,1930) demonstrated that an active principle could be obtained from the CL and the term progesterin

was used to designate this lipid-soluble substance. The structure of progesterone was later determined by Slotta et al. (1934). Subsequent research has sought to describe the hormonal control of CL function.

Formation of the CL is dependent on the morphological and biochemical transformation of follicular theca and granulosa cells into luteal cells (Gier and Marion, 1961; Donaldson and Hansel, 1965a). This transformation of follicular cells into luteal cells is dependent upon the preovulatory surge of the pituitary gonadotropins, luteinizing hormone (LH) and follicle-stimulating hormone (FSH) (reviewed by Channing et al., 1980). In the cow, the weight and progesterone (P_4) content of the CL increase rapidly between days 3 and 12 of the estrous cycle (estrus=day 0) and remain relatively constant until day 16 (Erb et al., 1971). At this time luteolysis, or regression of the CL, occurs followed by follicular recruitment, ovulation and subsequent CL development (reviewed by Hansel et al., 1973).

Two distinct populations of luteal cells have been identified in a variety of species such as the pig (Corner, 1919), cow (Foley and Greenstein, 1958), human (Gillim et al., 1969), sheep (Warbritton, 1934), rat (Wilkinson et al., 1976), white-tailed deer (Sinha et al., 1971), white whale (Mossman and Duke, 1973), and rhesus monkey (Guliyas et al., 1979). There is contradictory evidence on the follicular origin of these two cell types, designated small and large luteal cells. Several histological studies on the formation of the bovine CL suggest that luteal cells are derived from both cell types of the follicle, the granulosa giving rise to large luteal cells and the theca cells to the small luteal cells (McNutt, 1924; Foley and Greenstein, 1958; Gier and Marion, 1961; Donaldson and Hansel, 1965b). However, in the ewe and cow

It has been suggested that some of these large cells arise from small cells (Warbritton, 1934; Donaldson and Hansel, 1965b, Fitz et al., 1982). Recently, a study utilizing monoclonal antibodies directed against antigenic determinants of small and large luteal cells was conducted to determine the contribution of follicular cells to bovine luteal composition (Alla and Hansel, 1984). These authors reported that large cells were derived from granulosa while most of the small cells were of thecal origin, and that some small cells developed into large cells as the age of the CL increased. Additionally, they observed that granulosa-derived cells disappeared during early pregnancy while cells of thecal origin persisted throughout pregnancy.

Morphologically, large luteal cells are generally spherical in shape and their diameter greater than 20 μm , while small luteal cells have a stellate appearance and have a cell diameter of 10-20 μm (Priedkains and Weber, 1968; Koos and Hansel, 1981; Fitz et al., 1982). Ultrastructurally, both small and large luteal cells contain abundant endoplasmic reticulum and numerous mitochondria (Niswender et al., 1985); features which are characteristic of steroidogenically active cells (Christensen and Gillim, 1969; Savard, 1973). Large luteal cells also have features which are consistent with a protein secreting function, such as numerous Golgi complexes, rough endoplasmic reticulum and secretory granules (Niswender et al., 1985). These electron dense secretory granules are not characteristically found in small luteal cells although these cells do contain large numbers of lipid droplets (O'Shea et al., 1979; Koos and Hansel, 1981; Chengini et al., 1984; Rodgers et al., 1984; Niswender et al., 1985).

Although the follicular origin of small and large luteal cells is

controversial, the distinct steroidogenic capacities of these two cell types is more widely accepted. Differences in steroidogenic activity between small and large luteal cells was first reported by Leman and Loir (1977). They found that the basal rate of P_4 secretion by large cells was greater than that of small cells, and small cells were more responsive to LH. Similar results have now been reported utilizing luteal cell preparations from the cow (Ursely and Leymarie, 1979; Koos and Hansel, 1981), ewe (Fitz et al., 1982; Rodgers et al., 1983), and rabbit (Hoyer et al., 1986). Thus, both large and small cells secrete P_4 , but the regulation of steroidogenesis in the two cell types appears to be quite different. In the unstimulated state, large luteal cells secrete approximately 20 times as much P_4 as do small cells (Fitz et al., 1982). That LH can stimulate P_4 production in small cells is consistent with the finding that small cells contain more receptors for LH than do large cells (Fitz et al., 1982). In contrast, receptors for prostaglandins (Fitz et al., 1982) and estradiol (Glass et al., 1984) are more abundant on large cells. In addition to LH, steroidogenesis in small cells can be stimulated by dibutyryl cAMP or by agents that activate adenylate cyclase, such as cholera toxin and forskolin (Fitz et al., 1982; Rodgers et al., 1983; Hoyer et al., 1984). Therefore, steroidogenesis in small cells appears to be regulated through a cAMP-mediated mechanism. The mechanisms regulating steroid production in large luteal cells are not clear. Prostaglandins E_1 , E_2 and I_2 enhance P_4 production by large cells (Fitz et al., 1984a,b) while dibutyryl cAMP, cholera toxin and forskolin are ineffective in stimulating P_4 production by large cells (Fitz et al., 1982; Fitz et al., 1984a; Hoyer et al., 1984, 1986). These results suggest a mechanism of signal

transduction which is not associated with adenylate cyclase.

LUTEOTROPISM

A luteotropin is a substance which can maintain the morphological and biochemical integrity of the CL. The luteotropin varies among species and may consist of a single hormone or a complex of hormones. For example, prolactin (PRL), FSH and LH are required for luteal maintenance in the hamster (Greenwald, 1967, 1973) and mouse (Choredary and Greenwald, 1969; Hilliard, 1973) while PRL, LH and estradiol are required in the rat (Hilliard, 1973). In most cases the luteotropin is of pituitary origin; however, estrogen is luteotropic in some species such as the rabbit (Keyes and Naibandov, 1967) and rat (Hilliard, 1973). In the cow and ewe LH alone appears to be the major luteotropin. This conclusion is based upon a variety of studies utilizing in vivo and in vitro experimental models. In the cow and ewe corpora lutea either fail to form or regress following hypophysectomy at ovulation or mid-cycle, (Anderson et al., 1966; Kaltenbach et al., 1968a) and the effects of hypophysectomy can be reversed by the infusion of LH (Kaltenbach et al., 1968b). The in vivo administration of LH can overcome the luteolytic effects of oxytocin in the cow (Simmons and Hansel, 1964; Donaldson and Hansel, 1965b) and can prolong the lifespan of the CL (Wiltbank et al., 1961; Donaldson and Hansel, 1965b; Seguin et al., 1977). In hysterectomized heifers LH increased the total content and concentration of P_4 in the CL (Brunner et al., 1969), while administration of antisera to LH resulted in premature regression of the CL in the cow (Snook et al., 1969) and ewe (Fuller and Hansel, 1970). In vitro, the addition of

purified LH enhances the production of P_4 (Mason, 1962; Savard et al., 1965; Armstrong and Black, 1966; Selfart and Hansel, 1968), while the addition of LH antisera eliminates this response (Selfart and Hansel, 1968).

Although LH has received a considerable amount of attention as the primary luteotropin, an increasing body of literature has emerged concerning the role of other substances in the control of ovarian function. The following review of the literature concerns these other factors which may have a physiological role in the control of ovarian steroidogenesis.

FACTORS INFLUENCING OVARIAN FUNCTION

Catecholamines

The primary pathway catecholamines, norepinephrine (NE) and epinephrine (E), can stimulate the in vitro biosynthesis of cAMP and P_4 by luteal tissue from a variety of species: cow (Condon and Black, 1976; Godkin et al., 1977; Milvae and Hansel et al., 1983), rat (Harwood et al., 1979; Ratner et al., 1980; Norjavaara et al., 1982), rabbit (Birnbaumer et al., 1976), ewe (Jordan et al., 1978), and sow (Perkins et al., 1986). In addition the alternative pathway catecholamines octopamine, synephrine and deoxyepinephrine can also enhance P_4 production by bovine luteal cells (Battista and Condon, 1986b). In contrast to the stimulatory effects of catecholamines in the aforementioned species, the CL of the mare (Condon et al., 1979) and human (Richardson and Masson, 1984) are refractory to stimulation by exogenous adrenergic agonists.

In vivo, the infusion of NE, E or the β -adrenergic agonist isoproterenol (ISO) elevated plasma P_4 concentrations in the ewe (Bolt and Rollins, 1976), while infusion of NE increased cAMP levels in the CL of the rat (Norjavaara et al., 1983). In contrast, in vivo administration of NE and ISO was ineffective in stimulating ovarian steroidogenic activity in estrous rats, although fenoterol, a β_2 -adrenergic agonist, resulted in enhanced P_4 production (Zsolnai et al., 1982). Infusion of E in early pregnant women increased plasma P_4 (Fylling, 1971a), but these effects may have been due to stimulation of placental P_4 production, rather than a direct effect on luteal function (Fylling, 1971b; Flint et al., 1974; Csapo and Herczeg, 1977). This possibility is supported by the finding that human luteal tissue is unresponsive to catecholamines in vitro, as previously mentioned (Richardson and Masson, 1980; Casper and Cotterell, 1984). Infusion of ISO or the β -adrenergic antagonist propranolol in pseudopregnant rabbits did not alter luteal function (Gadsby et al., 1985). These authors suggested that endogenous catecholamines are not involved in the in vivo stimulation of luteal steroidogenesis. In the cow, administration of NE and E significantly elevated plasma P_4 concentrations, thus supporting a role for endogenous catecholamines in the control of bovine luteal function (Battista et al., 1987b).

Catecholamines act directly on luteal cells, since fibroblasts and erythrocytes separated from ewe luteal cell suspensions did not produce significant amounts of cAMP in response to E (Jordan et al., 1978). Adrenergic agents are thought to induce cAMP and subsequent P_4 production by binding to specific receptors in luteal cell membranes (Lefkowitz, 1976). Initial support for this hypothesis is suggested by

the finding that propranolol can block the catecholamine-induced increase in luteal cAMP and P_4 in the cow (Condon and Black, 1976; Godkin et al., 1978), rat (Harwood et al., 1980a; Norjavaara et al., 1982), and ewe (Jordan et al., 1978). These results not only suggest specific adrenergic receptors, but also that they are of the β -adrenergic type. Radioligand binding studies have demonstrated the presence of β -adrenergic receptors in rat luteal cell membranes (Coleman et al., 1978; Jordan, 1981; Kilachko and Zor, 1981). This receptor system has been further characterized to show that in the rat, the β -adrenergic stimulation of ovarian cAMP and P_4 is mediated by β_2 -adrenergic receptors (Ratner et al., 1980; Adashi and Hsueh, 1981a; Laszlovszky and Erdo, 1983; Norjavaara et al., 1984). In contrast, the CL of the cow (Battista and Condon, 1986a) and sow (Perkins et al., 1986) contains a mixed population of β -adrenergic receptors.

Serotonin and Melatonin

Serotonin (5-hydroxytryptamine; 5-HT) has been detected in whole ovarian homogenates of the rat (Clausell and Soliman, 1978) and in bovine luteal tissue (Battista et al., 1987a), while 3H -melatonin has been reported to be selectively taken up by the ovaries of the rat and cat in vivo (Wurtman et al., 1964). In vitro studies with human luteal tissue showed that melatonin stimulated P_4 production in a dose-dependent manner while 5-HT and N-acetyl 5-HT had no effect on P_4 production (MacPhee et al., 1975). In cultured rat granulosa cells, melatonin increased basal and LH-stimulated P_4 production (Fiske et al., 1984). In male rats melatonin and 5-HT inhibited the in vitro biosynthesis of androgens (Peat and Kinson, 1971; Ellis, 1972). In the cow, both 5-HT and 5-methoxytryptamine stimulate luteal P_4 secretion

while melatonin was ineffective in stimulating either basal or LH-stimulated P_4 production (Battista and Condon, 1986a; Battista et al., 1987a). Additionally, 5-HT has been shown to enhance bovine luteal P_4 production in vivo (Battista et al., 1987b).

Histamine and Dopamine

Although little is known about the role of histamine and dopamine in the control of ovarian function, Hirose et al. (1978) have reported that histamine stimulates cortisol and corticosterone production by dissociated adrenal cells of the dog. Greyard et al. (1983) found dopamine to inhibit the in vitro biosynthesis of androgens by rat testicular tissue. Utilizing bovine luteal tissue, Battista and Condon (1986a) were unable to demonstrate any stimulatory effect of histamine or a dopamine receptor agonist on basal or LH-stimulated P_4 production. However, we have recently found detectable levels of dopamine in bovine luteal tissue, and have found that dopamine stimulates P_4 production in vitro (Battista et al., unpublished observations).

Acetylcholine

As mentioned previously the mammalian ovary does receive cholinergic innervation, although the CL itself does not appear to contain cholinergic nerve fibers (Stefensen et al., 1981). That endogenous acetylcholine may play a role in influencing ovarian steroidogenesis is suggested by the findings that nicotinic, but not muscarinic agonists can inhibit FSH-induced P_4 production in cultured rat granulosa cells (Kasson and Hsueh, 1985a) and hCG-stimulated androgen biosynthesis in cultured rat testicular cells (Kasson and Hsueh, 1985b). In contrast, neither acetylcholine nor synthetic cholinergic agonists altered basal or LH-induced P_4 production by human luteal tissue in short- or long-

term culture (Casper and Cotterell, 1984). Similarly, bovine luteal tissue is unresponsive to acetylcholine (Battista and Condon, 1986a) or synthetic nicotinic and muscarinic agonists (Battista and Condon, unpublished observations).

Gamma-Aminobutyric Acid

A possible role for γ -aminobutyric acid (GABA) in the control of ovarian function has been suggested due to the relatively high concentration of this compound in rat ovarian tissue (Del Rio and Caballero, 1980; Erdo et al., 1982; Schaeffer and Hsueh, 1982; Apud et al., 1984). In addition glutamate decarboxylase, the enzyme responsible for the synthesis of GABA has been identified in whole ovarian homogenates but not in enriched ovarian granulosa cells (Schaeffer and Hsueh, 1982; Apud et al., 1984). Specific, high affinity GABA binding sites have been demonstrated in the rat and guinea pig ovary (Erdo and Lapis, 1982; Schaeffer and Hsueh, 1982; Amenta et al., 1986) with a majority of those sites being associated with granulosa cells (Schaeffer and Hsueh, 1982). These receptors have been further characterized to be of the GABA_A subtype (Amenta et al., 1986). In addition, a GABA-sensitive adenylyl cyclase (Erdo and Lapis, 1982) and high affinity sodium uptake system (Erdo, 1983) have been demonstrated in the rat ovary. With respect to ovarian function, GABA acts to increase ovarian blood flow and to enhance follicular estrogen secretion (Erdo et al., 1985). The role of GABA in the regulation of luteal steroidogenesis is unclear, although incubation of bovine luteal cells with GABA did not alter basal or LH-stimulated P₄ production (Battista and Condon, 1986a).

Vasoactive Intestinal Peptide

Immunoreactive vasoactive intestinal peptide (VIP), has been

detected within the ovarian stroma of the mature (Larsson et al., 1977; Alm et al., 1980a,b; Hakanson et al., 1982) and immature (Ahmed et al., 1986) rat ovary. VIP has been shown to stimulate, in a dose-dependent manner, the production of P_4 and estradiol in granulosa cells (Davoren and Hsueh, 1985) and whole ovarian homogenates (Ahmed et al., 1986) from the immature rat, and a role for VIP in the developmental regulation of ovarian steroidogenesis has been proposed (Ahmed et al., 1986). Intravenous infusion of VIP can elevate plasma P_4 concentrations in female rabbits (Fredricks et al., 1983), although this effect was later found to be due to VIP stimulation of adrenal P_4 secretion and this enhancement was independent of changes in LH and FSH (Fredricks et al., 1985). The role of VIP in the control of ovarian steroidogenesis in species other than the rat and rabbit is unknown.

Substance P

Nerve fibers containing substance P (SP) have been localized by immunofluorescence (Dees et al., 1985), radioimmunoassay and high performance liquid chromatography (Makris et al., 1982; Ojeda et al., 1985) in the immature rat ovary. The greatest amount of immunoreactive SP was associated with the ovarian vasculature, and no SP was detected in the CL (Dees et al., 1985). SP appears to be steroidogenically inactive since neither SP nor stable analogs altered basal or agonist-induced steroidogenesis in cultured granulosa cells or ovarian fragments obtained from immature rats (Ojeda et al., 1985). Whether SP is present in ovarian tissues and can modulate steroidogenesis in other species is unknown.

Opiates

Immunoreactive δ -endorphin has been localized in luteal tissue of

the rat (Tsong et al., 1982a; Lolait et al., 1985), mouse (Tsong et al., 1982a; Shana et al., 1984), and ewe (Lim et al., 1983), as well as in testicular cells of the rat and mouse (Tsong et al., 1982a,b). The concurrent detection of larger molecular weight species of immunoreactive β -endorphin as well as the detection of ACTH, MSH and proopiomelanocortin (POMC) is suggestive of de novo synthesis (Tsong et al., 1982b; Lim et al., 1983; Shana et al., 1984; Lolait et al., 1985), which appears to be regulated by gonadotropins (Shana et al., 1984; Lolait et al., 1985). A physiological role for opiate peptides in the control of luteal steroidogenesis has been suggested based on the observations of Abramowitz and Campbell (1983), who reported an enkephalin analog inhibited forskolin activation of adenylate cyclase in rabbit luteal membrane preparations. These observations suggest that the CL contains a receptor for opiate peptides which is coupled to an inhibitory guanine nucleotide component, and thus may serve as a negative autocrine regulator of luteal function (Abramowitz and Campbell, 1983). The role of opiate peptides in the control of steroidogenesis in other species has not been determined.

Oxytocin and Vasopressin

Oxytocin, oxytocin-associated neurophysin, and vasopressin have been observed in luteal tissue of the ewe (Wathes and Swann, 1982; Watkins 1983), cow (Fields et al., 1983; Wathes, 1983), pig (Wathes et al., 1983; Pitzel et al., 1984), and human (Wathes et al., 1982; Khan-Dawood and Dawood, 1983). The CL appears to be responsible for the synthesis of these peptides (Wathes and Swann, 1982; Wathes et al., 1982, 1983), and recently the expression of the oxytocin-specific mRNA was reported in bovine luteal tissue (Ivell et al., 1985).

The physiological role of these intraluteal peptides is not clear, however oxytocin has been implicated in the processes of luteolysis (Fairdough et al., 1980; McCracken, 1980; Flint and Sheldrick, 1983) and ovulation (Virutamasen et al., 1973, 1976).

With respect to regulating steroidogenesis, oxytocin and vasopressin have been reported to inhibit hCG-stimulated androgen synthesis in cultured rat Leydig cells (Adashi and Hsueh, 1981b). Tan et al. (1982a,b) have shown that high concentrations of oxytocin inhibited basal and hCG-enhanced P_4 production in bovine and human luteal cell preparations, while low concentrations of oxytocin enhanced basal steroidogenesis in both cell types. In contrast, oxytocin and vasopressin have no effect on basal or agonist-induced P_4 or cAMP production in luteal tissue of the rat (Mukhopadhyay et al., 1984) or human (Richardson and Masson, 1985). Similar results have been found for the effects of oxytocin in cultured bovine luteal cells (Burg, 1985).

Relaxin

The presence of relaxin has been demonstrated in luteal tissue of the cyclic and pregnant pig (Anderson et al., 1973a; Sherwood and O'Byrne, 1974; Sherwood and Rutherford, 1981), rat (Anderson et al., 1973b; Anderson et al., 1975; Anderson and Long, 1978; Sherwood and Rutherford, 1981), human (O'Byrne et al., 1978; Quagliarello et al., 1980), and pregnant cow (Fields et al., 1980, 1982). It is generally believed that the CL is the principle and perhaps sole source of relaxin in the pig, rat and cow, and the specific mRNA for relaxin has been identified in the porcine CL (Gast et al., 1980; Gast, 1982). In the human, the placenta may also serve as a source of relaxin (Fields and

Larkin, 1981). Although the factors directing the synthesis and secretion of relaxin have not been well characterized, the concentration of relaxin in the CL and peripheral circulation is elevated during the latter stages of gestation (reviewed by Sherwood and Downing, 1983). A physiological role for relaxin in the control of myometrial activity (Downing and Sherwood, 1985b) and parturition (Downing and Sherwood, 1985a,c) has been shown. A role for relaxin in the control of ovarian steroidogenesis has not been demonstrated.

Insulin and Insulin-Like Growth Factors

Insulin and the insulin-like growth factor somatomedin-C, have in recent years received considerable attention as regulators of steroidogenesis. Insulin enhanced basal and gonadotropin-induced P_4 production in rat and bovine luteal tissue (Tesone et al., 1983; Ladenheim et al., 1984; Poff et al., 1987), porcine granulosa cells (Channing et al., 1976; May et al., 1980; May and Schomberg, 1981; Veldhuis et al., 1983), and rat granulosa cells (Davoren and Hsueh, 1984; Adashi et al., 1984, 1985a,b). Somatomedin-C has similar steroidogenic activity in that it can synergize with FSH to enhance P_4 production in rat granulosa cells (Adashi et al., 1984; 1985a,b). The action of insulin and somatomedin-C is independent of cAMP (Adashi et al., 1984; Veldhuis and Furlanetto, 1985), while the phosphorylation of plasma membrane proteins (D'Erole et al., 1984) and the induction of components of the cholesterol side-chain cleavage system (Veldhuis et al., 1986) have been associated with them. Additionally, both insulin and somatomedin-C enhance the induction of LH-receptors in rat (Adashi et al., 1985b) and porcine (May et al., 1980; May and Schomberg, 1981) granulosa cells. These observations, plus the finding of discrete

binding sites for insulin and possibly for somatomedin-C (Hsueh et al., 1984; Ladenheim et al., 1984; Adashi et al., 1985a,b), and the observation that granulosa cells are capable of synthesizing somatomedin-like peptides (Hammond et al., 1983, 1984; Adashi et al., 1985b) suggest an autocrine role for insulin and insulin-like growth factors in the control of ovarian function.

Epidermal Growth Factor

Specific binding sites for epidermal growth factor (EGF) have been demonstrated in luteal, thecal and granulosa cells of the rat (Jones et al., 1982; Arnaud et al., 1983; Chabot et al., 1983) and cow (Vlodavsky et al., 1978). In the rat, EGF receptor concentration varies with the stage of the estrous cycle, with the highest concentration occurring at estrus (Arnaud et al., 1983). Additionally, EGF receptor synthesis was found to be regulated by gonadotropins (Arnaud et al., 1983).

With respect to steroidogenesis, EGF inhibited FSH-induced estrogen production in cultured rat granulosa cells (Hsueh et al., 1981; Welsh and Hsueh, 1982), and the production of P_4 in cultured porcine (Channing et al., 1983) and bovine (Savion and Gospodarowicz, 1980; Franchimont et al., 1986) granulosa cells. In contrast, EGF enhanced basal and agonist-induced P_4 production in granulosa cells of the rat (Jones et al., 1982; Knecht and Catt, 1983). The divergent effects of EGF in regulation of rat granulosa cell steroidogenesis is associated with EGF inhibition of aromatase activity and stimulation of 3β -hydroxysteroid dehydrogenase activity (Jones et al., 1982).

Gonadotropin-Releasing Hormone

Treatment with gonadotropin-releasing hormone (GnRH) or its agonists in vivo results in paradoxical inhibition of female reproductive

function (Corbin et al., 1978; Nilius et al., 1978; Rivier et al., 1978). In vitro, GnRH or its analogues inhibited gonadotropin-induced cAMP and steroidogenesis in rat granulosa (Hsueh and Erickson, 1979; Hsueh and Ling, 1979; Jones and Hsueh, 1981) and luteal cells (Clayton et al., 1979; Behrman et al., 1980; Harwood et al., 1980b). Furthermore, specific GnRH receptor sites have been identified in rat follicular and luteal tissue (Clayton et al., 1979; Harwood et al., 1980b; Pieper et al., 1981). In addition, the inhibitory effects of GnRH on granulosa cells involved the activation of a plasma membrane cAMP phosphodiesterase (Knecht and Catt, 1981). Treatment of rat granulosa cells with GnRH alone enhanced cAMP and progesterone production (Knecht et al., 1981; Clark et al., 1982; Jones and Hsueh, 1982a,b; Ranta et al., 1982), although these effects were not observed in luteal cells (Behrman et al., 1980; Harwood et al., 1980).

In contrast to the effects of GnRH on steroidogenesis in the rat, no direct effects of GnRH have been observed in ovarian tissues of the human (Tan and Biggs, 1983), monkey (Asch et al., 1981a), or cow (Kuman, 1984). In addition no GnRH binding was observed in ovarian tissues of the human (Clayton, 1982) or monkey (Asch et al., 1981b).

Adenosine

Adenosine and other adenine-derived purines can amplify LH-dependent cAMP accumulation and P_4 production in rat (Hall et al., 1981; Behrman et al., 1982; Brennan et al., 1983) and human (Polan et al., 1983) luteal tissue. Adenosine can also enhance FSH-dependent cAMP and P_4 production in human and rat granulosa cells (Polan et al., 1983). The stimulatory effects of adenosine have also been shown in rat Leydig and adrenal cells (Wolff and Hope-Cook, 1977; Rommerts et al., 1984).

Moreover, the antigonadotropic effect of $\text{PGF}_2\alpha$ on luteal cAMP and P_4 production was completely antagonized by adenosine (Behrman et al., 1982).

The amplification of gonadotropin-induced cAMP accumulation by adenosine is dependent upon an intracellular site of action since the effects of adenosine can be blocked by dipyridamol, an inhibitor of nucleoside transport (Hall et al., 1981; Brennan et al., 1983), and a selective transport system for adenosine has been demonstrated in rat luteal cells (Behrman et al., 1983). The mechanism underlying adenosine amplification of LH-dependent cAMP accumulation is due to utilization of extracellular adenosine as a substrate for ATP formation, and thus acts as a prosubstrate for conversion into cAMP by adenylate cyclase (Behrman et al., 1983; Brennan et al., 1983). Whether an adenosine uptake mechanism is functional in bovine luteal cells is at this time unknown.

Prostaglandins

The bovine CL can synthesize prostaglandins, and this synthesis can be stimulated by LH (Shemesh and Hansel, 1975a,b; Hixon and Hansel, 1975; Milvae and Hansel, 1983) and inhibited by indomethacin (Milvae and Hansel, 1985). Rat granulosa cells can also synthesize prostaglandins in vitro, and LH can act to increase prostaglandin synthetase activity (Clark et al., 1978; Koos and Clark, 1982). Prostaglandin receptors have been identified in a variety of species including the cow (Kimball and Lauderdale, 1975; Rao, 1974,1975), ewe (Powell et al., 1974a), rabbit (Abramowitz and Birnbaumer, 1979), and human (Powell et al., 1974b).

Prostaglandins of the E-series (PGE_1 and PGE_2) and I-series (PGI_2) interact with luteal membrane receptors to enhance cAMP and P_4

production (Marsh, 1971; Abramowitz and Birnbaumer, 1979; Milvae and Hansel, 1980, 1983). While prostaglandins of the F-series (notably PGF_2^α) act to inhibit luteal steroidogenesis in vitro (for review of literature see Pate and Condon, 1984) and in vivo (reviewed by Inskeep and Murdoch, 1980) and a physiological role for PGF_2^α in luteolysis (McCracken et al., 1972; McCracken, 1980) and ovulation (reviewed by Espey, 1978) has been proposed.

Glucocorticoids

Administration of pharmacological doses of glucocorticoids blocked ovulation in women and laboratory animals (Hagino, 1972). Treatment of rat granulosa cells with cortisol, inhibited FSH-induced estrogen production (Hsueh and Erickson, 1978), while stimulating the production of P_4 (Adashi et al., 1981). The divergent effects of glucocorticoids on rat granulosa cell steroidogenesis is associated with inhibition of aromatase activity and stimulation of 3β -hydroxysteroid dehydrogenase activity (Adashi et al., 1981). In male rats, in vivo treatment with glucocorticoids decreased plasma testosterone concentrations (Saez et al., 1977; Bambino and Hsueh, 1976, 1979). Incubation of cultured rat Leydig cells with glucocorticoids inhibited basal and LH-induced androgen production and reduced LH-receptor synthesis (Bambino and Hsueh, 1981).

That glucocorticoids may be of physiological significance in the control of steroidogenesis is suggested by the presence of receptors in rat granulosa cells (Lorevel et al., 1977) and in rat testicular tissue (Ballard et al., 1974; Evain et al., 1976). Data concerning the effects of glucocorticoid treatment on luteal steroidogenesis is not available.

Catecholestrogens

The catecholestrogens 2- and 4-hydroxyestradiol are naturally occurring compounds formed from estradiol by the enzyme estrogen-2/4-hydroxylase, and are considered to be the major biproducts of estradiol metabolism in mammals (reviewed by Ball and Knuppen, 1980). Catecholestrogens can mimic or alter estrogen responses through their interaction with estrogen receptors (MacLusky et al., 1983), and can influence enzymes regulating the biosynthesis and catabolism of catecholamines (Lloyd and Welsz, 1978).

The role of estradiol in the enhancement of gonadotropin-induced steroidogenesis and receptor induction in rat ovarian tissue is well established (reviewed by Hsueh et al., 1984). Recently, catecholestrogens have been shown to enhance steroidogenesis in rat granulosa (Hudson and Hillier, 1985) and luteal cells (Khan and Gabori, 1984). These results, plus the observation of catecholesterogen formation in rat (Barbieri et al., 1978) and porcine (Hammond et al., 1986) follicular tissue, suggest that catecholestrogens are intraovarian regulators of steroidogenesis.

Despite the existence of estrogen receptors in rat follicular and luteal tissue (Richards, 1974), it does not appear that catecholestrogens are acting as weak estrogens (Hudson and Hillier, 1985; Hammond et al., 1986). Recently Barnea and Fakh (1985) reported that catecholestrogens stimulate placental steroidogenesis, and that their effects were mediated by α and β -adrenergic receptors. Thus, it is possible that catecholestrogens may enhance steroidogenesis through interactions with β -adrenergic receptors, which are present on rat ovarian cells (Coleman et al., 1979). Both the cow and ewe have luteal

estradiol receptors (Kimball and Hansel, 1974; Glass et al., 1984) as well as β -adrenergic receptors (Condon and Black, 1976; Jordan et al., 1978). However, estradiol is luteolytic in both of these species (reviewed by Hansel, 1973; Williams and Marsh, 1978). The possibility that catecholestrogens may enhance luteal steroidogenesis in the cow and ewe through interaction with β -adrenergic receptors has not been investigated.

MECHANISMS CONTROLLING STEROIDOGENESIS

Hormone-Receptor Interaction: Activation of Adenylate Cyclase

Initial events mediating steroidogenesis involve the interaction of a hormone with a specific membrane receptor. As discussed in the previous section, luteal cells (and/or granulosa cells) contain a variety of receptors which mediate both stimulatory and inhibitory control on steroidogenesis (reviewed by Hsueh et al., 1983;1984). Hormone-receptor interaction results in the formation of a hormone-receptor complex (H-R complex) which interacts with a regulatory protein which then activates adenylate cyclase, a membrane-associated enzyme which catalyzes the conversion of MgATP to cAMP (Marsh, 1970). Other membrane-associated events which are coupled to H-R interaction include membrane methylation, which results in alteration of membrane fluidity (Hirata et al., 1979; Milvae et al., 1983). The decrease in membrane microviscosity acts to increase the lateral mobility of the H-R complex, thus enhancing coupling to adenylate cyclase and unmasking membrane receptors (reviewed by Hirata and Axelrod, 1980; Axelrod and Hirata, 1982). Additionally, H-R interaction is associated with clustering and

Internalization of membrane receptors (reviewed by Willingham and Pastan, 1984), which are either packaged into vesicles for degradation (Ahmed et al., 1981) or remain biologically active and further alter intracellular responses (Zimniski et al., 1982; Rao et al., 1983).

The hormone-sensitive adenylate cyclase system is composed of three membrane bound components; a receptor which has a high affinity and specificity for the ligand, a guanine nucleotide regulatory protein or G-protein (N), and a catalytic unit (reviewed by Rodbell, 1980; Birnbaumer and Kirchick, 1983). The regulatory unit is a dimeric protein composed of two subunits designated N_{α} and N_{β} . The N_{α} contains the binding sites for GTP and the catalytic unit, while N_{β} contains a Mg ion binding site and the ability to interact with the H-R complex (reviewed by Hildebrand et al., 1984a). Upon binding of the hormone to its receptor the H-R complex couples with N_{β} and causes a shift in the K_m of N for Mg from 10mM to less than 0.5mM, which is the physiological concentration of Mg. Binding of Mg to the N_{β} portion induces dissociation of the N protein allowing N_{α} to interact with and activate the catalytic unit. There is also a requirement for the nucleotide GTP, which when bound to N_{α} allows it to couple with the catalytic unit. The activated catalytic unit is then able to enhance the conversion of MgATP to cAMP. Therefore, hormones act primarily by reducing the Mg requirement for GTP-dependent activation of adenylate cyclase. Return of the system to the nonstimulated state is dependent upon the GTPase activity associated with N_{α} . Thus hydrolysis of GTP reverses the binding of N_{α} to the catalytic unit resulting in inactivation of adenylate cyclase (reviewed by Hildebrand et al., 1984; Birnbaumer et al., 1985a,b). The inhibitory effects of various hormones and

neurotransmitters upon adenylate cyclase and subsequent cAMP formation are mediated by a distinct GTP regulatory protein designated N_1 (Jakobs, 1979; Cooper, 1982; Gilman, 1984). N_1 , like N_5 , undergoes subunit dissociation with subsequent binding to and inactivation of the catalytic unit. The exact mechanism whereby N_1 attenuates the activity of the catalytic unit is unknown (reviewed by Rodbell, 1980; Birnbaumer et al., 1985a,b).

cAMP and cAMP-Dependent Protein Kinase

Addition of cAMP to luteal tissue increased P_4 production while incubation with LH resulted in a rapid accumulation of cAMP prior to P_4 synthesis. Subsequently it was determined that the action of LH was mediated by adenylate cyclase rather than the inhibition of phosphodiesterase activity (reviewed by Marsh, 1976; Williams et al., 1978). The biological effects of cAMP are mediated through a specific cAMP-dependent protein kinase (PK_A), the activation of which involved agonist-induced steroidogenesis (Menon, 1973; Vaitukaitis et al., 1975; Azhar et al., 1976; Ling and Marsh, 1977). PK_A is composed of regulatory and catalytic subunits, which in the inactive state exist as a holoenzyme. Binding of two molecules of cAMP to the regulatory units induces subunit dissociation and release of two enzymatically active catalytic subunits (reviewed by Walsh and Ashley, 1973; Krebs and Beavo, 1979). The active catalytic subunit can then phosphorylate a variety of intracellular proteins, and a correlation in protein phosphorylation and P_4 synthesis has been observed in luteal tissue (Darbon et al., 1981).

Regulation of Steroidogenesis: Protein Kinase A

There are three sources of cholesterol which can be utilized for the synthesis of P_4 : de novo synthesis from acetate, cholesterol derived

from circulating lipoproteins, or cholesterol derived from stored cholesterol esters (reviewed by Dorrington, 1977). The de novo synthesis of cholesterol is regulated by the enzyme 3-hydroxy-3-methylglutaryl-coenzyme A reductase (Brown et al., 1973), which is regulated by a phosphorylation-dephosphorylation mechanism (Beg et al., 1980; Ingebritsen et al., 1981), the activity of which is induced by LH (Kovanen et al., 1978; Schuler et al., 1981). PK_A can also phosphorylate and activate cholesterol ester hydrolase resulting in the hydrolysis of cholesterol esters to yield free cholesterol (Trzeciak and Boyd, 1973; Beckett and Boyd, 1977; Cooke et al., 1983; Colbran et al., 1986). Intracellular cholesterol must then be transported to the mitochondria where it is converted to pregnenolone in a rate limiting reaction catalyzed by the cholesterol side-chain cleavage enzyme system (reviewed by Savard, 1973). The side-chain cleavage system resides in or on the inner mitochondrial membrane and is composed of three proteins: 1) luteodoxin reductase, an FAD-containing flavoprotein, 2) cytochrome $P_{450_{SCC}}$, the terminal oxygenase, and 3) luteodoxin, which serves to shuttle electrons between the reductase and cP_{450} (reviewed by Sulimovici and Boyd, 1969). The cholesterol side-chain cleavage reaction involves two sequential hydroxylations of cholesterol at carbon 22 and carbon 20 followed by cleavage of the cholesterol side-chain between carbon 20 and 22. Each step in the reaction requires oxygen and NADPH, thus the conversion of one mole of cholesterol to one mole of pregnenolone requires three moles of oxygen and NADPH (reviewed by Burstein and Gut, 1971).

Many investigators have demonstrated that agonist-induced steroidogenesis is dependent upon protein synthesis (reviewed by Garren

et al., 1971; Savard et al., 1976). Although it has been suggested that PK_A acts directly on the side-chain cleavage system (Caron et al., 1975; Williams et al., 1978), recent reports have shown that PK_A has no direct regulatory action on mitochondrial steroidogenesis (Downing and Dimino, 1979; Inaba and Wiest, 1985). The observation that the transport of cholesterol to the inner mitochondrial membrane is translation dependent (Crivello and Jefcoate, 1980; Mori and Marsh, 1982), coupled with the finding that PK_A is translocated to the nucleus where it may regulate transcription (Jungmann et al., 1974; Spielvogel et al., 1977), lead to the proposal that stimulation of steroidogenesis is dependent upon the synthesis of a labile regulatory protein (Simpson et al., 1978). The labile protein is thought to be generated on cytoplasmic ribosomes, since chloramphenicol, an inhibitor of mitochondrial protein synthesis, was ineffective in blocking steroidogenesis (Arthur and Boyd, 1974). The labile protein may enhance agonist-induced steroidogenesis by aiding in the transport of cholesterol across the mitochondrial membrane, thereby enhancing cholesterol side-chain cleavage activity (reviewed by Strauss et al., 1981).

After the conversion of cholesterol to pregnenolone, pregnenolone diffuses out of the mitochondria to the smooth endoplasmic reticulum where it is converted to P_4 by the enzymes $\Delta^5-3\beta$ -hydroxysteroid dehydrogenase (3β -HSD) and $\Delta^5 \Delta^4$ -isomerase (Tamaoki, 1973). The enzyme 20α -hydroxysteroid dehydrogenase (20α -HSD) is also located in the smooth endoplasmic reticulum and acts to convert P_4 to the less potent progestin 20β -dihydroprogesterone (Tamaoki, 1973). With respect to the control of steroidogenesis, both 3β -HSD and 20α -HSD are under hormonal regulation in rat granulosa cells. Stimulatory agents such as LH, FSH

catecholamines increase the enzymatic activity of 3 β -HSD while decreasing 20 α -HSD activity. Inhibitory substances, such as GnRH, act to regulate these enzymes in a reverse fashion. The mechanism whereby these enzymes are regulated in other species is unknown (reviewed by Hsueh et al., 1983, 1984).

Regulation of Steroidogenesis: Protein Kinase C

Protein kinase C (PK_C) is a cytosolic enzyme composed of a single polypeptide chain. The enzyme does not appear to consist of subunits since activation is not associated with dissociation of the enzyme molecule (reviewed by Kuo et al., 1984). Activation of PK_C is dependent upon phosphatidylinositol (PI) hydrolysis and intracellular Ca²⁺ (reviewed by Kuo et al., 1984; Nishizuka, 1984). PI comprises less than ten percent of the total phospholipid in mammalian membranes. Most of it is associated with the plasma membrane although some is associated with the mitochondria and the endoplasmic reticulum (reviewed by Mitchell, 1975). Hydrolysis of PI is associated with agonist-induced phosphorylation of PI by specific kinases to form phosphatidylinositol 4-phosphate (PIP) and phosphatidylinositol 4,5-bisphosphate (PIP₂). PIP₂ is acted upon by phospholipase-C to yield diacylglycerol (DAG) and inositol triphosphate (IP₃). IP₃ acts to enhance the intracellular mobilization of Ca²⁺, thus promoting the production of DAG by the Ca²⁺-dependent phospholipase-C (reviewed by Farese, 1983a,b, 1984). DAG, in conjunction with Ca²⁺, activates PK_C by increasing its affinity for Ca²⁺. The activated PK_C then acts synergistically with mobilized Ca²⁺ to induce cellular responses. Thus, PK_C is a Ca²⁺-dependent, phospholipid-sensitive enzyme, which when activated acts in a synergistic manner with intracellular Ca²⁺ to elicit a full physiological response (reviewed by

Kuo et al., 1984; Nishizuka, 1984).

With respect to steroidogenesis, LH has been shown to increase PI hydrolysis in bovine luteal tissue (Davis et al., 1981) and rat granulosa cells (Davis et al., 1983). PK_C activity has been demonstrated in cytosolic preparations from the bovine CL (Davis and Clark, 1983) and human ovary (Clark et al., 1983). Use of phorbol esters, which can substitute for DAG and directly activate PK_C (Castagna et al., 1982; Niedel et al., 1983), have been shown to stimulate steroidogenesis in bovine luteal tissue (Brunswig et al., 1986) and rat granulosa cells (Kawai and Clark, 1985; Shinohara et al., 1986). The mechanism whereby PK_C mediates agonist-induced steroidogenesis is unknown.

Progesterone Secretion

Once synthesized, P_4 must be secreted from the CL into the bloodstream. It has been generally accepted over the years that the synthesis of steroid hormones is directly coupled to stimulation, and that newly synthesized steroids leave the cells within minutes by passive diffusion (Vogt, 1943; Short, 1964; Enders, 1973; Baird, 1977; Rice et al., 1986). However, some data suggest that the secretion of P_4 may be an active process. For example, P_4 secretion requires Ca^{2+} (Higuchi et al., 1976), and K^+ -induced membrane depolarization is associated with a rapid enhancement in P_4 secretion (Higuchi et al., 1976). It has been suggested that P_4 binds to cytosolic proteins and is packaged into granules which are released by an exocytotic mechanism (Gemmell et al., 1974; Quirk et al., 1979; Sawyer et al., 1979). Two specific P_4 binding proteins have been found in cytosolic preparations of bovine luteal tissue (Willcox, 1983). However, alteration of

cytoskeletal structures important in cellular transport were found to have no effect on P_4 secretion (Zor et al., 1978; Azhar and Menon, 1981; Gwynne and Condon, 1982). Additionally, Rice et al. (1986) were unable to demonstrate the presence of P_4 in secretory granules within the ovine corpus luteum. Furthermore, evidence suggests that instead of secretory granules, serum steroid binding proteins may facilitate the transport of steroid hormones from their site of synthesis into the blood stream (Condon and Pate, 1981).

REFERENCES

- Abramowitz, J. and Birnbaumer, L. (1979). Prostacyclin activation of adenylyl cyclase in rabbit corpus luteum membranes: comparison with 6-keto-prostaglandin $F_{1\alpha}$ and E_1 . *Biol. Reprod.* 21:609-616.
- Abramowitz, J. and Campbell, A.R. (1983). Enkephalin-mediated inhibition of forskolin-stimulated rabbit luteal adenylyl cyclase activity. *Biochem. Biophys. Res. Comm.* 116:574-580.
- Adashi, E.Y. and Hsueh, A.J.W. (1981a). Stimulation of β_2 -adrenergic responsiveness by follicle-stimulating hormone in rat granulosa cells *in vitro* and *in vivo*. *Endocrinology* 108:2170-2178.
- Adashi, E.Y. and Hsueh, A.J.W. (1981b). Direct inhibition of testicular androgen biosynthesis revealing antigonadal activity of neurohypophyseal hormones. *Nature* 293:650-652.
- Adashi, E.Y., Jones, P.B.C. and Hsueh, A.J.W. (1981). Synergistic effect of glucocorticoids on the stimulation of progesterone production by follicle-stimulating hormone in cultured rat granulosa cells. *Endocrinology* 109:1888-1894.
- Adashi, E.Y., Resnick, C.E., Svoboda, M.E. and Van Wyk, J.J. (1984). A novel role for somatomedin-C in the cytodifferentiation of the ovarian granulosa cell. *Endocrinology* 115:1227-1229.
- Adashi, E.Y., Resnick, C.E., Svoboda, M.E. and Van Wyk, J.J. (1985a). Somatomedin-C synergizes with follicle-stimulating hormone in the acquisition of progestin biosynthetic capacity by cultured rat granulosa cells. *Endocrinology* 116:2135-2142.
- Adashi, E.Y., Resnick, C.E., Svoboda, M.E. and Van Wyk, J.J. (1985b). Somatomedin-C enhances induction of luteinizing hormone receptors by follicle-stimulating hormone in cultured rat granulosa cells. *Endocrinology* 116:2369-2375.
- Aguado, L.I., Petrovic, S.L. and Ojeda, S.R. (1982). Ovarian β -adrenergic receptors during the onset of puberty: characterization, distribution and coupling to steroidogenic responses. *Endocrinology* 111:1124-1131.
- Aguado, L.I. and Ojeda, S.R. (1984a). Prepubertal ovarian function is finely regulated by direct adrenergic influences. Role of noradrenergic innervation. *Endocrinology* 114:1845-1853.
- Aguado, L.I. and Ojeda, S.R. (1984b). Ovarian adrenergic nerves play a role in maintaining preovulatory steroid secretion. *Endocrinology* 114:1944-1946.

- Ahmed, C.E., Sawyer, H.R. and Niswender, G.D. (1981). Internalization and degradation of human chorionic gonadotropin in ovine luteal cells: kinetic studies. *Endocrinology* 109:1380-1387.
- Ahmed, C.E., Dees, W.L. and Ojeda, S.R. (1986). The immature rat ovary is innervated by vasoactive intestinal peptide (VIP)-containing fibers and responds to VIP with steroid secretion. *Endocrinology* 118:1682-1689.
- Alla, H.W. and Hansel, W. (1984). Origin of different cell types in the bovine corpus luteum as characterized by specific monoclonal antibodies. *Biol. Reprod.* 31:1015-1025.
- Allen, W.M. and Corner, G.W. (1930). Physiology of corpus luteum VII. Maintenance of pregnancy in rabbit after very early castration, by corpus luteum extracts. *Proc. Soc. Exp. Biol. Med.* 27:403-405.
- Alm, P., Alumets, J., Hakanson, R., Owman, C., Sjöberg, N.O., Sundler, F. and Wallis, B. (1980a). Origin and distribution of VIP (vasoactive intestinal peptide) nerves in the genito-urinary tract. *Cell Tissue Res.* 205:337-347.
- Alm, P., Alumets, J., Hakanson, R., Owman, C., Sjöberg, N.O., Sundler, F. and Wallis, B. (1980b). Vasoactive intestinal peptide nerves in the human female genital tract. *Am. J. Obstet. Gynecol.* 136:349-351.
- Alm, P., Alumets, J., Hakanson, R., Owman, Ch., Sjöberg, N.O., Sternquist, M. and Sundler, F. (1983). Enkephalin-immunoreactive nerve fibers in the feline genito-urinary tract. *Histochem.* 72:351-355.
- Amenta, F., Endo, S.L., Mione, M.C. and Napoleone, P. (1986). ³H-Muscimol binding sites within guinea pig ovary: a histoautoradiographic study. *Pharmacol.* 32:202-207.
- Amoroso, E.C. and Perry, J.S. (1977). Ovarian activity during gestation. In: *The Ovary*, 2nd edn., Vol. II, Physiology. (S. Zuckerman and B.J. Weir, eds.). Academic Press, New York, pp.315-398.
- Anderson, L.L., Oxenreider, S.L., Hendricks, D.M. and Melampy, R.M. (1966). Pituitary stalk section in beef heifers. *J. Anim. Sci.* 25:1262-1263.
- Anderson, L.L., Ford, J.J., Melampy, R.M. and Cox, D.F. (1973a). Relaxin in porcine corpora lutea during pregnancy and after hysterectomy. *Am. J. Physiol.* 225:1215-1219.
- Anderson, L.L., Blast, J.P. and Melampy, R.M. (1973b). Relaxin in ovarian tissue during different reproductive states in the rat. *J. Endocrinol.* 59:371-372.
- Anderson, M.L., Long, J.A. and Hayashida, T. (1975). Immunofluorescence studies on the localization of relaxin in the corpus luteum of the

- pregnant rat. Biol. Reprod. 13:499-504.
- Anderson, M.L. and Long, J.A. (1978). Localization of relaxin in the pregnant rat. Bioassay of tissue extracts and cell fractionation studies. Biol.Reprod. 18:110-117.
- Apud, J.A., Tappaz, M.L., Celotti, F., Negrì-Cesi, P., Masotto, C. and Racagni, G. (1984). Biochemical and immunochemical studies on the GABAergic system in the rat fallopian tube and ovary. J. Neurochem. 43:120-125.
- Armstrong, D.T. and Black, D.L. (1966). Influence of luteinizing hormone on corpus luteum metabolism and progesterone biosynthesis throughout the bovine estrous cycle. Endocrinology 78:937-944.
- Arnaud, R.st., Walker, P., Kelly, P.A. and Labrie, F. (1983). Hormonal regulation of ovarian epidermal growth factor (EGF) receptors in the rat. In: Factors Regulating Ovarian Function. (G.S. Greenwald and P.F. Terranova, eds.). Raven Press, New York, pp.209-213.
- Arthur, J.R. and Boyd, G.S. (1974). The effect of inhibitors of protein synthesis on cholesterol side chain-cleavage in the mitochondria of luteinized rat ovaries. Eur. J. Biochem. 49:117-127.
- Asch, R.H., Eddy, C.A. and Schally, A.V. (1981a). Lack of luteolytic effect of D-TRP-6-LH-RH in hypophysectomized rhesus monkeys (*Macaca mulatta*). Biol. Reprod. 25:963-968.
- Asch, R.H., Sickle, M.V., Rettori, V., Balmaceda, J.P., Eddy, C.A., Coy, D.H., and Schally, A.V. (1981b). Absence of LH-RH binding sites in corpora lutea from rhesus monkeys (*Macacca mulatta*). J. Clin. Endocrinol. Metab. 53:215-217.
- Axelrod, J. and Hirata, F. (1982). Phospholipid methylation and the receptor-induced release of histamine from cells. TIPS 3:156-158.
- Azhar, S., Clark, M.R. and Menon, K.M.J. (1976). Regulation of cyclic adenosine 3',5'-monophosphate dependent protein kinase of rat ovarian cells by luteinizing hormone and human chorionic gonadotropin. Endocrine Res. Comm. 3:93-104.
- Azhar, S. and Menon, K.M.J. (1981). Receptor-mediated gonadotropin action in the ovary. Action of cytoskeletal element-disrupting agents on gonadotropin-induced steroidogenesis in rat luteal cells. Biochem. J. 194:19-27.
- Bahr, J., Kao, L. and Naibandov, A.V. (1974). The role of catecholamines and nerves in ovulation. Biol. Reprod. 10:273-290.
- Bahr, J.M. and Ben-Johnathan, N. (1981). Preovulatory depletion of ovarian catecholamines in the rat. Endocrinology 108:1815-1820.
- Bahr, J.M. and Ben-Johnathan, N. (1985). Elevated catecholamines in porcine follicular fluid before ovulation. Endocrinology 117:620-623.

- Baird, D.T. (1977). Synthesis and secretion of steroid hormones by the ovary. In: *The Ovary, Vol. 3, Regulation of Oogenesis and Steroidogenesis*. (L. Zuckerman and B.J. Weir, eds.). Academic Press, New York, pp. 305-357.
- Ball, P. and Knuppen, R. (1980). Catecholamines: Chemistry, biogenesis, metabolism, occurrence and physiological significance. *Acta Endocrinol.* 232:1-127.
- Ballard, P.D., Baxter, J.D., Higgins, S.J., Rousseau, G.G., and Tomkins, G.M. (1974). General presence of glucocorticoid receptors in mammalian tissues. *Endocrinology* 94:998-1002.
- Bambino, T.H. and Hsueh, A.J.W. (1981). Direct inhibitory effect of glucocorticoids upon testicular luteinizing hormone receptor and steroidogenesis *in vivo* and *in vitro*. *Endocrinology* 108:2142-2148.
- Barbieri, R.L., Canick, J.A. and Ryan, K.J. (1978). Estrogen 2-hydroxylase: activity in rat tissues. *Steroids* 32:529-538.
- Barnea, E.R. and Fakh, H.N. (1985). The role of catecholestrogens in placental steroidogenesis. *Steroids* 45:427-432.
- Battista, P.J. and Condon, W.A. (1986a). Serotonin-induced stimulation of progesterone production by cow luteal cells *in vitro*. *J. Reprod. Fert.* 76:231-238.
- Battista, P.J. and Condon, W.A. (1986b). A role for alternative pathway catecholamines in the regulation of steroidogenesis in cow luteal cells. *J. Reprod. Fert.* 78:275-280.
- Battista, P.J., Rexroad, C.E. Jr. and Condon, W.A. (1987a). Mechanisms involved in the action of serotonin-induced stimulation of progesterone production by bovine luteal cells *in vitro*. *Molec. Cellular Endocrinol.*, In press.
- Battista, P.J., Poff, J.P., Deaver, D.R. and Condon, W.A. (1987b). Biogenic amine regulation of bovine luteal progesterone production *in vivo*. *J. Reprod. Fert.*, In press.
- Beckett, G.J. and Boyd, G.S. (1977). Purification and control of bovine adrenal cortical cholesterol ester hydrolase and evidence for the activation of the enzyme by a phosphorylation. *Eur. J. Biochem.* 72:223-233.
- Beg, Z.H., Stonik, J.A. and Brewer, H.B. (1980). *In vitro* and *in vivo* phosphorylation of rat liver 3-hydroxy-3-methylglutaryl coenzyme A reductase and its modulation by glucagon. *J. Biol. Chem.* 255:8541-8545.
- Behrman, H.R., Preston, S.L. and Hall, A.K. (1980). Cellular mechanism of the antigonadotropic action of luteinizing hormone releasing hormone in the corpus luteum. *Endocrinology* 107:656-664.

- Behrman, H.R., Hall, A.K., Preston, S.L. and Gore, S.D. (1982). Antagonistic interactions of adenosine and prostaglandin F_2^{α} modulate acute responses of luteal cells to luteinizing hormone. *Endocrinology* 110:38-46.
- Behrman, H.R., Ohkwawa, R., Preston, S.L. and MacDonald, G.J. (1983). Transport and selective utilization of adenosine as a prosubstrate for luteinizing hormone-sensitive adenylate cyclase in the luteal cell. *Endocrinology* 113:1132-1140.
- Bell, C. (1972). Autonomic control of reproduction. *Pharmacol. Rev.* 24:657-736.
- Ben-Johnathan, N., Arbogast, L.A., Rhodes, T.A. and Bahr, J. (1984). Norepinephrine in the rat ovary: Ontogeny and *de novo* synthesis. *Endocrinology* 115:1426-1431.
- Birnbaumer, L., Yang, P.C., Hunzicker-Dunn, M., Brockaert, J. and Duran, J.M. (1976). Adenyl cyclase activities in ovarian tissues. I. Homogenization and conditions of assay in Graafian follicles and corpora lutea of rabbits, rats and pigs: regulation by ATP, and some comparative properties. *Endocrinology* 99:163-184.
- Birnbaumer, L. and Kirchick, H.J. (1983). Regulation of gonadotropic action: The molecular mechanisms of gonadotropin-induced activation of ovarian adenyl cyclase. In: *Factors Regulating Ovarian Function*. (G.S. Greenwald and P.F. Terranova, eds.). Raven Press, New York, pp. 287-310.
- Birnbaumer, L., Codina, J., Mattera, R., Carlone, R.A., Hildebrandt, J.D., Sunyer, T., Rojas, F.J., Caron, M.G., Lefkowitz, R.J. and Iyengar, R. (1985a). Structural basis of adenylate cyclase stimulation and inhibition by distinct guanine nucleotide regulatory proteins. In: *Molecular Mechanisms of Transmembrane Signaling*. (Cohn and Houslay, eds.). Elsevier Science Publishers, New York. pp. 131-182.
- Birnbaumer, L., Codina, J., Mattera, R., Carlone, R.A., Hildebrandt, J.D., Sunyer, T., Rojas, F.J., Caron, M.G., Lefkowitz, R.J. and Iyengar, R. (1985b). Regulation of hormone receptors and adenyl cyclases by guanine nucleotide binding N proteins. *Rec. Prog. Horm. Res.* 41:41-99.
- Black, D.L. (1974). Neural control of oviduct musculature. In: *The Oviduct and its Functions*. (A.D. Johnson and C.W. Foley, eds.). Academic Press, New York, pp. 65-118.
- Black, D.L. (1979). Catecholamines-Influences on corpora lutea and tubal genitalia. In: *Beltville Symposia in Agricultural Research, Animal Reproduction*. 3:373-397.
- Blank, M.A., Gu, J., Allen, J.M., Haung, W.M., Yiangou, Y., Ch'ng, J., Lewis, G., Elder, M.G., Polak, J.M. and Bloom, S.R. (1986). The regional distribution of NPY-, PHM-, and VIP-containing nerves in the human female genital tract. *Int. J. Fert.* 31:218-222.

- Bolt, D.J. and Rollins, R. (1976). Increased plasma progesterone during infusion of epinephrine, norepinephrine or isoproterenol in luteal phase ewes. *Biol. Reprod. Suppl.* 1:43 (abstract).
- Brennan, T.J., Ohkawa, R., Gore, S.D. and Behrman, H.R. (1983). Adenosine-derived purines increase ATP levels in the luteal cell: evidence that cell levels of ATP may limit cAMP accumulation by luteinizing hormone. *Endocrinology* 112:499-508.
- Brink, C.E. and Grob, H.S. (1972). Response of the denervated mouse ovary to exogenous gonadotropins. *Biol. Reprod.* 9:108 (abstract).
- Brown, M.S., Dana, S.E. and Goldstein, J.L. (1973). Regulation of 3-hydroxy-3-methylglutaryl coenzyme A reductase activity in human fibroblast by lipoproteins. *Proc. Natl. Acad. Sci.* 70:2162-2166.
- Brundin, J. (1965). Distribution and function of adrenergic nerves in the rabbit fallopian tube. *Acta Physiol. Scand.* 66:suppl. 259:1-20.
- Brunner, M.A., Donaldson, L.E. and Hansel, W. (1969). Exogenous hormones and luteal function in hysterectomized and intact helpers. *J. Dairy Sci.* 52:1849-1854.
- Brunswig, B., Mukhopadhyay, A.K., Budnik, L.T., Bohnet, H.G. and Leidenberger, F.A. (1986). Phorbol ester stimulates progesterone production by isolated bovine luteal cells. *Endocrinology* 118:743-749.
- Burden, H.W. (1972). Adrenergic innervation in the ovaries of the rat and guinea pig. *Am. J. Anat.* 133:455-462.
- Burden, H.W. (1978). Ovarian innervation. In: *The Vertebrate Ovary*. (R.E. Jones, ed.). Plenum Press, New York, pp. 615-638.
- Burden, H.W. and Lawrence, L.E. (1977). The effects of denervation on the localization of $\Delta^5-3\beta$ -hydroxysteroid dehydrogenase activity in the rat ovary during pregnancy. *Acta Anat.* 97:286-290.
- Burden, H.W. and Lawrence, L.E. (1978). Experimental studies on the acetylcholinesterase-positive nerves in the ovary of the rat. *Anat. Rec.* 190:233-242.
- Burg, M.F. (1985). Comparison of differential responses of early and midcycle bovine luteal cells to hormonal treatments in a serum-free cell culture system. Masters Thesis, University of New Hampshire.
- Burstein, S. and Gut, M. (1971). Biosynthesis of pregnenolone. *Rec. Prog. Horm. Res.* 27:303-349.
- Carlson, R.R. and DeFeo, V.J. (1965). Role of the pelvic nerve vs. the abdominal sympathetic nerves in the reproductive function of the female rat. *Endocrinology* 77:1014-1022.
- Caron, M.C., Goldstein, S., Savard, K. and Marsh, J.M. (1975). Protein

- kinase stimulation of a reconstituted cholesterol side-chain cleavage enzyme system in the bovine corpus luteum. *J. Biol. Chem.* 250:5137-5143.
- Casper, R.F. and Cotterell, M.A. (1984). The effects of adrenergic and cholinergic agents on progesterone production by human corpus luteum in vitro. *Am. J. Obstet. Gynecol.* 148:663-669.
- Castagna, M., Takai, Y., Kaibuchi, K., Sano, K., Kikkawa, U. and Nishizuka, Y. (1982). Direct activation of calcium-activated, phospholipid-dependent protein kinase by tumor-promoting phorbol ester. *J. Biol. Chem.* 257:7847-7851.
- Chabot, J.G., Arnaud, R. St., Walker, P. and Pelletier, G. (1986). Distribution of epidermal growth factor receptors in the rat ovary. *Molec. Cellular Endocrinol.* 44:99-108.
- Channing, C.P. and Kammerman, S. (1974). Binding of gonadotropins to ovarian cells. *Biol. Reprod.* 10:179-188.
- Channing, C.P., Tsai, V. and Sachs, D. (1976). Role of insulin, thyroxine and cortisol in luteinization of porcine granulosa cells grown in chemically defined media. *Biol. Reprod.* 15:235-247.
- Channing, C.P., Schaert, F.W., Anderson, L.D. and Tsafiriri, A. (1980). Ovarian follicular and luteal physiology. In: *Reproductive Physiology III, International Review of Physiology*, Vol. 22. (R.O. Greep, ed.). University Park Press, Baltimore, pp. 117-201.
- Channing, C.P., Garrett, R., Kroman, N., Conn, T. and Gospodarowicz, D. (1983). Ability of EGF and FGF to cause a decrease in progesterone secretion by cultured porcine granulosa cells: changes in responsiveness throughout follicular maturation. In: *Factors Regulating Ovarian Function*. (G.S. Greenwald and F.F. Terranova, eds.). Raven Press, New York, pp. 215-220.
- Chegini, N., Ramani, N. and Rao, Ch. V. (1984). Morphological and biochemical characterization of small and large bovine luteal cells during pregnancy. *Molec. Cellular Endocrinol.* 37:89-102.
- Christensen, A.K. and Gillim, S.W. (1969). The correlation of fine structure and function in steroid-secreting cells, with emphasis on those of the gonads. In: *The Gonads*. (K.W. McKerns, ed.). Appleton, New York, pp. 415-488.
- Choudary, J.B. and Greenwald, G.S. (1969). Luteotropic complex in the mouse. *Anat. Rec.* 163:373-387.
- Clark, M.R., Marsh, J.M. and LeMaire, W.J. (1978). Mechanism of luteinizing hormone regulation of prostaglandin synthesis in rat granulosa cells. *J. Biol. Chem.* 253:7757-7761.
- Clark, M.R. (1982). Stimulation of progesterone and prostaglandin E accumulation by luteinizing-hormone releasing hormone (LHRH) and LHRH analogs in rat granulosa cells. *Endocrinology* 110:146-152.

- Clark, M.R., Davis, J.S. and LeMaire, W.J. (1983). Calcium- and lipid-dependent protein phosphorylation in the human ovary. *J. Clin. Endocrinol. Metab.* 57:872-875.
- Clausell, D.E. and Soliman, K.F.A. (1978). Ovarian serotonin content in relation to ovulation. *Experientia* 34:410-411.
- Clayton, R.N., Harwood, J.P. and Catt, K.J. (1979). Gonadotropin-releasing hormone analogue binds to luteal cells and inhibits progesterone production. *Nature* 282:90-92.
- Clayton, R.N. (1982). Absence of gonadotropin-releasing hormone receptors in human gonadal tissue. *Nature* 299:56-59.
- Colbran, R.J., Garton, A.J., Cordle, S.R. and Yeaman, S.J. (1986). Regulation of cholesterol ester hydrolase by cyclic AMP-dependent protein kinase. *FEBS* 201:257-261.
- Coleman, A.J., Paterson, D.S. and Somerville, A.R. (1979). The adrenergic receptor of rat corpus luteum membranes. *Biochem. Pharmacol.* 28:1003-1010.
- Condon, W.A. and Black, D.L. (1976). Catecholamine-induced stimulation of progesterone by the bovine corpus luteum *in vitro*. *Biol. Reprod.* 15:573-578.
- Condon, W.A., Ganjam, V.K. and Kenney, R.M. (1979). Catecholamines and equine luteal progestins. *J. Reprod. Fert., Suppl.* 27:199-203.
- Condon, W.A. and Pate, J.L. (1981). Influence of serum and its lipoprotein fractions on progesterone synthesis and secretion by bovine luteal tissue *in vitro*. *Biol. Reprod.* 25:950-957.
- Cook, K.G., Colbran, R.J., Snee, J. and Yeaman, S.J. (1983). Cytosolic cholesterol ester hydrolase from bovine corpus luteum its purification, identification, and relationship to hormone-sensitive lipase. *Biochim. Biophys. Acta* 752:46-53.
- Corbin, A., Beattie, C.W., Tracy, J., Foell, T.J., Yardley, J. and Rees, R.W.A. (1978). The anti-reproductive pharmacology of LH-RH and agonistic analogues. *Int. J. Fert.* 23:81-92.
- Corner, G.W. (1919). On the origin of the corpus luteum of the sow from both granulosa and theca interna. *Am. J. Anat.* 26:117-183.
- Corner, G.W. and Allen, W.M. (1929). Physiology of corpus luteum II. Production of a specific uterine reaction (progestinal proliferation) by extract of the corpus luteum. *Am. J. Physiol.* 88:326-339.
- Crievello, J.F. and Jefcoate, C.R. (1980). Intracellular movement of cholesterol in rat adrenal cells, kinetics and effects of inhibitors. *J. Biol. Chem.* 255:8144-8151.

- Curry, T.E., Lawrence, I.E. and Burden, H.W. (1983). Ovarian follicular development following exogenous gonadotropin stimulation in the prepubertal guinea pig: effect of unilateral 6-hydroxydopamine. *Fed. Proc.* 42:316 (abstract).
- Darbon, J.M., Ursely, J. and Leymarie, P. (1981). Correlation between protein phosphorylation and progesterone synthesis in bovine luteal cells stimulated by lutropin. *Eur. J. Biochem.* 119:237-243.
- Davoren, J.B. and Hsueh, A.J.W. (1984). Insulin enhances FSH-stimulated steroidogenesis by cultured rat granulosa cells. *Molec. Cellular Endocrinol.* 35:97-105.
- Davoren, J.B. and Hsueh, A.J.W. (1985). Vasoactive intestinal peptide: a novel stimulator of steroidogenesis by cultured rat granulosa cells. *Biol. Reprod.* 33:37-52.
- Davis, J.S., Farese, R.V. and Marsh, J.M. (1981). Stimulation of phospholipid labeling and steroidogenesis by luteinizing hormone in isolated bovine luteal cells. *Endocrinology* 109:469-475.
- Davis, J.S., Farese, R.V. and Clark, M.R. (1983). Stimulation of phospholipid synthesis by luteinizing hormone in isolated rat granulosa cells. *Endocrinology* 112:2212-2214.
- Davis, J.S. and Clark, M.R. (1983). Activation of protein kinase in the bovine corpus luteum by phospholipids and Ca^{2+} . *Biochem. J.* 214:569-574.
- Dees, W.L., Kozlowski, Dey, R. and Ojeda, S.R. (1985). Evidence for the existence of substance P in the prepubertal rat ovary. II. Immunocytochemical localization. *Biol. Reprod.* 33:471-476.
- Del Rio, M.C. (1981). Gamma-aminobutyric acid system in rat oviduct. *J. Biol. Chem.* 256:9816-9819.
- Del Rio, M.C. and Caballero, A.L. (1980). Presence of γ -aminobutyric acid in rat ovary. *J. Neurochem.* 34:1584-1586.
- D'Ercole, A.J., Stiles, A.D. and Underwood, L.E. (1984). Tissue concentrations of somatomedin-C: further evidence for multiple sites of synthesis and paracrine or autocrine mechanisms of action. *Proc. Natl. Acad. Sci.* 81:935-939.
- Doerr, P. and Pirke, K.M. (1976). Cortisol induced suppression of plasma testosterone in normal adult males. *J. Clin. Endocrinol. Metab.* 43:622-629.
- Doerr, P. and Pirke, K.M. (1979). Dexamethasone induced suppression of the circadian rhythm of plasma testosterone in normal adult males. *J. Steroid Biochem.* 10:81-86.
- Donaldson, L. and Hansel, W. (1965a). Histological study of the bovine corpus luteum. *J. Dairy Sci.* 48:905-909.

- Donaldson, L.E. and Hansel, W. (1965b). Prolongation of the lifespan of the bovine corpus luteum by single injections of bovine luteinizing hormone. *J. Dairy Sci.* 48:903-904.
- Dorrington, J.H. (1977). Steroidogenesis in vitro. In: *The Ovary, Vol.3, Regulation of Oogenesis and Steroidogenesis*. (S. Zuckerman and B.J. Weir, eds.). Academic Press, New York, pp. 359-412.
- Downing, J.R. and Dimino, M.J. (1979). Studies on mitochondrial protein kinase activity by porcine corpora lutea. *Endocrinology* 105:570-573.
- Downing, S.J. and Sherwood, O.D. (1985a). The physiological role of relaxin in the pregnant rat. I. The influence of relaxin on parturition. *Endocrinology* 116:1200-1205.
- Downing, S.J. and Sherwood, O.D. (1985b). The physiological role of relaxin in the pregnant rat. II. The influence of relaxin on uterine contractile activity. *Endocrinology* 116:1206-1214.
- Downing, S.J. and Sherwood, O.D. (1985c). The physiological role of relaxin in the pregnant rat. III. The influence of relaxin on cervical extensibility. *Endocrinology* 116:1215-1220.
- Eik-Nes, K.B. (1969). An effect of isoproterenol on rates of synthesis and secretion of testosterone. *Am. J. Physiol.* 217:1764-1770.
- Ellis, L.C. (1972). Inhibition of rat testicular androgen synthesis in vitro by melatonin and serotonin. *Endocrinology* 90:17-28.
- Enders, A.C. (1973). Cytology of the corpus luteum. *Biol. Reprod.* 8:158-182.
- Erb, R.E., Randel, R.D. and Callahan, C.J. (1971). Female sex steroid changes during the reproductive cycle. *J. Anim. Sci.* suppl. 32:80-106.
- Erdo, S.L. (1983). High affinity sodium dependent aminobutyric acid uptake by slices of rat ovary. *J. Neurochem.* 40:582-594.
- Erdo, S.L. and Lapis, E. (1982). Bicuculline-sensitive GABA receptors in the rat ovary. *Eur. J. Pharmacol.* 85:243-246.
- Erdo, S.L., Rosdy, B. and Szporny, L. (1982). Higher GABA concentrations in follicular tube than in brain of the rat. *J. Neurochem.* 38:1174-1176.
- Erdo, S.L., Varga, B. and Horvath, E. (1985). Effect of local GABA administration on rat ovarian blood flow and progesterone and estradiol secretion. *Eur. J. Pharmacol.* 111:397-400.
- Espey, L.L. (1978). Ovulation. In: *The Vertebrate Ovary*. (R.E. Jones, ed.). Plenum Press, New York, pp. 503-532.
- Evain, D., Morera, A.M. and Saez, J.M. (1976). Glucocorticoid receptors in interstitial cells of the rat testis. *J. Steroid Biochem.* 7:1135-

1139.

- Fairclough, R.J., Moore, L.G., McGowam, L.T., Peterson, A.J., Smith, J.F., Tervit, H.R. and Watkins, W.B. (1980). Temporal relationship between plasma concentrations of 13,14-dihydro-15-keto-prostaglandin F and neurophysin I/II around luteolysis in sheep. *Prostaglandins* 20:199-208.
- Farese, R.V. (1983). The role of the phosphatidate-insolite cycle in the action of steroidogenic agents. *J. Steroid Biochem.* 19:1029-1032.
- Farese, R.V. (1984). Phospholipids as intermediates in hormone action. *Molec. Cellular Endocrinol.* 35:1-14.
- Farese, R.V. (1983). Phosphoinositide metabolism and hormone action. *Endocr. Rev.* 4:78-95.
- Farrar, J.A., Handberg, G.M., Hartley, M.L. and Pennefather, J.N. (1980). Catecholamine levels in the guinea pig ovary, myometrium and costo-uterine muscle during the estrous cycle and in the ovary after unilateral ovariectomy. *Biol. Reprod.* 22:473-479.
- Fields, M.J., Fields, P.A., Castro-Hernandez, A. and Larkin, L.H. (1980). Evidence for relaxin in corpora lutea of late pregnant cows. *Endocrinology* 107:869-876.
- Fields, M.J., Roberts, R. and Fields, P.A. (1982). Octadecylsilica and carboxymethyl cellulose isolation of bovine and porcine relaxin. *Ann. NY Acad. Sci.* 380:36-46.
- Fields, P.A. and Larkin, L.H. (1981). Purification and immunohistochemical localization of relaxin in the human term placenta. *J. Clin. Endocrinol. Metab.* 52:79-85.
- Fields, P.A., Eldridge, R.K., Fuchs, A.R., Roberts, R.F. and Fields, M.J. (1983). Human placental and bovine corpora luteal oxytocin. *Endocrinology* 112:1544-1546.
- Fiske, V.M., Parker, K.L., Ulmer, R.A., Ow, C.H. and Aziz, N. (1984). Effects of melatonin alone or in combination with human chorionic gonadotropin or ovine luteinizing hormone on the *in vitro* secretion of estrogens or progesterone by granulosa cells of rats. *Endocrinology* 114:407-410.
- Fitz, T.A., Mayan, M.H., Sawyer, H.R. and Niswender, G.D. (1982). Characterization of two steroidogenic cell types in the ovine corpus luteum. *Biol. Reprod.* 27:703-711.
- Fitz, T.A., Hoyer, P.B. and Niswender, G.D. (1984a). Interactions of prostaglandins with subpopulations of ovine luteal cells. I. Stimulatory effects of prostaglandins E₁, E₂ and I₂. *Prostaglandins* 28:119-126.
- Fitz, T.A., Mock, E.J., Mayan, M.H. and Niswender, G.D. (1984b).

Interactions of prostaglandins with subpopulations of ovine luteal cells. II. Inhibitory effects of $\text{PGF}_2\alpha$ and protection by PGE_2 . *Prostaglandins* 28:127-138.

Flint, A.P.F., Anderson, A.B.M. and Turnbull, A.C. (1974). Control of utero-ovarian and peripheral venous plasma progesterone by β -sympathomimetic drugs in pregnant sheep. *J. Endocrinol.* 63:253-254.

Flint, A.P.F. and Sheldrick, E.L. (1983). Ovarian secretion of oxytocin is stimulated by prostaglandin. *Nature* 297:587-588.

Foley, R.C. and Greenstein, J.S. (1958). Cytological changes in the bovine corpus luteum during early pregnancy. In: *Reproduction and Fertility*. (F.X. Gassner, ed.). Pergamon Press, New York, pp.88-96.

Ford, S.P. (1982). Control of uterine and ovarian blood flow throughout the estrous cycle and pregnancy of ewes, sows and cows. *J. Anim. Sci. Suppl.* 2:32-42.

Foster, D.L., Yellon, S.M. and Oister, D.H. (1985). Internal and external determinants of the timing of puberty in the female. *J. Reprod. Fert.* 75:327-344.

Fraenkel, L. and Cohn, F. (1901). Experimentelle Untersuchungen über den Einfluss des Corpus Luteum auf die Insertion des Eies. *Anat. Anz.* 20:294-300.

Franchimont, P., Hazee-Hagelstein, M.T., Charlet-Renard, Ch. and Jaspar, J.M. (1986). Effect of mouse epidermal growth factor on DNA and protein syntheses, progesterone and inhibin production by bovine granulosa cells in culture. *Acta Endocrinol.* 111:122-127.

Fredricks, C.M., Lundquist, L.E., Mathur, R.S., Ashton, S.H. and Landgrebe, S.C. (1983). Effects of vasoactive intestinal polypeptide upon ovarian steroids, ovum transport and fertility in the rabbit. *Biol. Reprod.* 28:1052-1060.

Fredricks, C.M., Ashton, S.H., Anderson, W.F., Mathur, R.S., Lundquist, L.E. and Landgrebe, S. (1985). Vasoactive intestinal polypeptide stimulates nonovarian progesterone secretion in rabbits. *Peptides* 6:205-210.

Fuller, G.B. and Hansel, W. (1970). Regression of sheep corpora lutea after treatment with antiovine luteinizing hormone. *J. Anim. Sci.* 31:99-103.

Fylling, P. (1971a). Vasopressin-induced increase in peripheral plasma progesterone levels during early human pregnancy. *Acta Endocrinol. Copenh.* 66:273-282.

Fylling, P. (1971b). Propranolol-blockade of vasopressin induced increase in plasma progesterone in early human pregnancy. *Acta Endocrinol. Copenh.* 66:283-288.

Gadsby, J.E., Keyes, P.L., Schwartz, T.S., Bill, C.H. and Lucchesi, B.

- (1985). Do catecholamines play a physiological role in regulating corpus luteum function in the pseudopregnant rabbit? *Biol. Reprod.* 32:907-915.
- Garren, L.D., Gill, G.N., Masul, H. and Walton, G.M. (1971). On the mechanism of action of ACTH. *Rec Prog. Horm. Res.* 27:433-478.
- Gelder, J.B. and Pepler, R.D. (1979). Effect of spinal cord transection on the reproductive system in the female rat. *Neuroendocrinol.* 29:293-298.
- Gemmell, R.T., Stacy, B.D. and Thorburn, G.D. (1974). Ultrastructural study of secretory granules in the corpus luteum of the sheep during the estrous cycle. *Biol. Reprod.* 11:447-462.
- Gerendel, I., Marchetti, B., Maugen, S., Amico-Roxas, M. and Scapagnini, V. (1978). Prevention of compensatory ovarian hypertrophy by local treatment of the ovarian hypertrophy by local treatment of the ovary hypertrophy by local treatment of the ovary with 6-OHDA. *Neuroendocrinol.* 27:272-280.
- Gier, H.T. and Marlon, G.B. (1961). The formation of the bovine corpus luteum. *J. Dairy Sci.* 44:1187.
- Gillim, S.W., Christensen, A.K. and McLennan, C.E. (1969). Fine structure of the human menstrual corpus luteum at its stage of maximum secretory activity. *Am. J. Anat.* 126:409-428.
- Gilman, A.G. (1984). G proteins and dual control of adenylate cyclase. *Cell* 36:577-579.
- Glass, J.D., Filtz, T.A. and Niswender, G.D. (1984). Cytosolic receptor for estradiol in the corpus luteum of the ewe: Variation throughout the estrous cycle and distribution between large and small steroidogenic cell types. *Biol. Reprod.* 31:967-974.
- Godkin, J.D., Black, D.L. and Duby, R.T. (1977). Stimulation of cyclic AMP and progesterone synthesis by LH, PGE₂ and isoproterenol in the bovine CL *in vitro*. *Biol. Reprod.* 17:514-518.
- Greenwald, G.S. (1967). Luteotropic complex of the hamster. *Endocrinology* 80:118-130.
- Greenwald, G.S. (1973). Further evidence for a luteotropic complex in the hamster: progesterone determinations of plasma and corpora lutea. *Endocrinology* 92:235-242.
- Grizard, G., Andre, M., Jarrige, J.F., Chambon, M. and Boucher, D. (1983). Effects of bromocriptine on pituitary-testicular function in the rat: possible inhibition of *in vitro* production of androgen by leydig cells. *Int. J. Androl.* 6:563-568.
- Grob, H.S. (1974). Follicular growth in denervated ovaries of hypophysectomized mice. *Fed. Proc.* 33:213 (abstract).

- Gulyas, B.J., Stouffer, R.L. and Hodgen, G.P. (1979). Progesterone synthesis and fine structure of dissociated monkey (*Macaca mulatta*) luteal cells maintained in culture. *Biol. Reprod.* 20:779-792.
- Gwynne, A. and Condon, W.A. (1982). Effects of cytochalasin B, Colchicine, and vinblastine on progesterone synthesis and secretion by bovine luteal tissue in vitro. *J. Reprod. Fert.* 65:151-156.
- Hagino, N. (1972). The effect of synthetic glucocorticoids on ovarian function in the baboon. *J. Clin. Endocrinol. Metab.* 35:716-721.
- Hakanson, R., Sundler, F. and Uddman, R. (1982). Distribution and topography of peripheral VIP nerve fibers: functional implications. In: *Vasoactive Intestinal Peptide*. (S.I. Said, ed.). Raven Press, New York, pp.121-132.
- Hall, A.K., Preston, S.L. and Behrman, H.R. (1981). Purine amplification of luteinizing hormone action in ovarian luteal cells. *J. Biol. Chem.* 256:10390-10398.
- Hammond, J.M., Yoshida, K., Veldhuis, J.D., Rechler, M.M. and Knight, A.P. (1983). Intrafollicular role of somatomedins: comparison with effects of insulin. In: *Factors Regulating Ovarian Function*. (G.S. Greenwald and P.F. Terranova, eds.). Raven Press, New York, pp. 197-201.
- Hammond, J.M., Baranao, J.L.F., Skaleris, D.A., Rechler, M.M. and Knight, A.P. (1984). Somatomedin (Sm) production by cultured porcine granulosa cells (GC). *J. Steroid Biochem.* 20:1597.
- Hammond, J.M., Hersey, R.M., Walega, M.A. and Welsz, J. (1986). Catecholestrogen production by porcine ovarian cells. *Endocrinology* 118:2292-2299.
- Hansel, W., Concannon, P.W. and Lukaszewska, J.H. (1973). Corpora lutea of large domestic animals. *Biol. Reprod.* 8:222-245.
- Harwood, J.P., Dufau, M.L. and Catt, K.J. (1979). Differing specificities of ovarian cyclase by epinephrine and human chorionic gonadotropin. *Molec. Pharmacol.* 15:439-444.
- Harwood, J.P., Richert, N.D., Dufau, M.L. and Catt, K.J. (1980a). Gonadotropin-induced desensitization of epinephrine action in the luteinized rat ovary. *Endocrinology* 107:280-288.
- Harwood, J.P., Clayton, R.N. and Catt, K.J. (1980b). Ovarian gonadotropin-releasing hormone receptors. I. Properties and inhibition of luteal function. *Endocrinology* 107:407-413.
- Higuchi, T., Kaneko, A., Abel, J.H. Jr. and Niswender, G.D. (1976). Relationship between membrane potential and progesterone release in ovine corpora lutea. *Endocrinology* 99:1023-1032.
- Hildebrandt, J.D., Codina, J., Iyengar, R., Rojas, F.J., Kirchick, H.J., Abromowitz, J., Hunzicker-Dunn, M. and Birnbaumer, L. (1984a).

- Transmembrane regulation of adenylyl cyclases. In: *Hormone Receptors in Growth and Reproduction*. (B.B. Saxena, ed.). Raven Press, New York, pp.111-140.
- Hildebrandt, J.D., Codina, J., Risper, R. and Birnbaumer, L. (1984b). Identification of a γ -subunit associated with adenylyl cyclase regulatory proteins N_5 and N_1 . *J. Biol. Chem.* 259:2039-2042.
- Hill, R.T. (1949). Adrenal cortical physiology of spleen grafted and denervated ovaries in the mouse. *Exp. Med. Surg.* 7:86-98.
- Hill, R.T. (1962). Paradoxical effects of ovarian secretions. In: *The Ovary*, Vol. 2. (S. Zuckerman, ed.). Academic Press, New York, pp.231-261.
- Hilliard, J. (1973). Corpus luteum function in guinea pigs, hamsters, rats, mice and rabbits. *Biol. Reprod.* 8:203-221.
- Hirata, F., Strittmatter, W.J. and Axelrod, J. (1979). β -Adrenergic receptor agonists increase phospholipid methylation, membrane fluidity, and β -adrenergic receptor-adenylate cyclase coupling. *Proc. Natl. Acad. Sci.* 76:368-372.
- Hirata, F. and Axelrod, J. (1980). Phospholipid methylation and biological signal transmission. *Science* 209:1082-1090.
- Hirose, T., Matsumoto, I. and Aikawa, T. (1978). Direct effect of histamine on cortisol and corticosterone production by isolated dog adrenal cells. *J. Endocrinol.* 76:371-372.
- Hixon, J.E. and Hansel, W. (1979). Effects of prostaglandin F_2^{α} , estradiol and luteinizing hormone in dispersed cell preparations of bovine corpora lutea. In: *Ovarian Follicular and Corpus Luteum Function*. (C.P. Channing, J.M. Marsh, W.A. Sadler, eds.). Plenum Press, New York, pp.613-620.
- Hoyer, P.B., Fitz, T.A. and Niswender, G.D. (1984). Hormone independent activation of adenylyl cyclase in large steroidogenic ovine luteal cells does not result in increased progesterone secretion. *Endocrinology* 114:604-608.
- Hoyer, P.B., Keyes, P.L. and Niswender, G.D. (1986). Size distribution and hormonal responsiveness of dispersed rabbit luteal cells during pseudopregnancy. *Biol. Reprod.* 34:905-910.
- Hoyer, P.B. and Niswender, G.D. (1986). Adenosine 3',5'-monophosphate-binding capacity in small and large ovine luteal cells. *Endocrinology* 119:1822-1829.
- Hsueh, A.J.W. and Erickson, G.F. (1978). Glucocorticoid-inhibition of FSH-induced estrogen production in cultured rat granulosa cells. *Steroids* 32:639-648.
- Hsueh, A.J.W. and Erickson, G.F. (1979). Extrapituitary action of gonadotropin-releasing hormone: Direct inhibition of luteal

steroidogenesis. *Science* 204:854-855.

- Hsueh, A.J.W. and Ling, N.C. (1979). Effect of an antagonistic analog of gonadotropin releasing hormone upon ovarian granulosa cell function. *Life Sci.* 25:1223-1230.
- Hsueh, A.J.W., Welsh, T.H. and Jones, P.B.C. (1981). Inhibition of ovarian and testicular steroidogenesis by epidermal growth factor. *Endocrinology* 108:2002-2004.
- Hsueh, A.J.W., Jones, P.B.C., Adashi, E.Y., Wang, C., Zhuang, L.Z. and Welsh, T.H. Jr. (1983). Intraovarian mechanisms in the hormonal control of granulosa cell differentiation in rats. *J. Reprod. Fert.* 69:325-342.
- Hsueh, A.J.W., Adashi, E.Y., Jones, P.B.C. and Welsh, T.H. (1984). Hormonal regulation of the differentiation of cultured ovarian granulosa cells. *Endocr. Rev.* 5:76-91.
- Hudson, K.E. and Hillier, S.G. (1985). Catechol oestradiol control of FSH-stimulated granulosa cell steroidogenesis. *J. Endocrinol.* 106:R1-R4.
- Inaba, T. and Wiest, W.A. (1986). Protein kinase stimulation of steroidogenesis in rat luteal mitochondria. *Endocrinology* 117:315-322.
- Ingelblitsen, T.S., Parker, R.A. and Gibson, D.M. (1981). Regulation of liver hydroxymethylglutaryl-CoA reductase by a bicyclic phosphorylation system. *J. Biol. Chem.* 256:1138-1144.
- Inskeep, E.K. and Murdoch, W.J. (1980). Relation of ovarian functions to uterine and ovarian secretion of prostaglandins during the estrous cycle and early pregnancy in the ewe and cow. In: *International Review of Physiology*. (R.O. Greep, ed.), University Park Press, Baltimore, pp.325-357.
- Ivell, R., Brackett, K.H., Fields, M.J. and Richter, D. (1985). Ovulation triggers oxytocin gene expression in the bovine ovary. *FEBS Lett.* 190:263-267.
- Jacobowitz, D. and Wallach, E.E. (1967). Histochemical and chemical studies of the autonomic innervation of the ovary. *Endocrinology* 81:1132-1139.
- Jakobs, K.H. (1979). Inhibition of adenylate cyclase by hormones and neurotransmitters. *Mol. Cell. Endocrinol.* 16:147-156.
- Jones, P.B.C. and Hsueh, A.J.W. (1981). Direct effects of gonadotropin releasing hormone and its antagonists upon ovarian functions stimulated by FSH, prolactin and LH. *Biol. Reprod.* 24:747-759.
- Jones, P.B.C. and Hsueh, A.J.W. (1982a). Regulation of ovarian 3 - hydroxysteroid dehydrogenase by gonadotropin-releasing hormone and follicle-stimulating hormone in cultured rat granulosa cells.

Endocrinology 110:1663-1671.

- Jones, P.B.C. and Hsueh, A.J.W. (1982b). Pregnenolone biosynthesis by cultured rat granulosa cells: modulation by follicle-stimulating hormone and gonadotropin-releasing hormone. *Endocrinology* 111:713-721.
- Jones, P.B.C., Welsh, T.H. and Hsueh, A.J.W. (1982). Regulation of ovarian progesterone production by epidermal growth factor in cultured rat granulosa cells. *J. Biol. Chem.* 257:11268-11273.
- Jordan, A.W. III, Caffrey, J.L. and Niswender, G.D. (1978). Catecholamine-induced stimulation of progesterone and adenosine 3', 5'-monophosphate production by dispersed ovine luteal cells. *Endocrinology* 103:385-392.
- Jungmann, R.A., Hiestand, P.C. and Schueppe, J.S. (1974). Adenosine 3',5'-monophosphate-dependent protein kinase and the stimulation of ovarian nuclear ribonucleic acid polymerase activities. *J. Biol. Chem.* 249:5444-5451.
- Kaltenbach, C.C., Graber, J.W., Niswender, G.D. and Naibandov, A.V. (1968a). Effect of hypophysectomy on the formation and maintenance of corpora lutea in the ewe. *Endocrinology* 82:753-759.
- Kaltenbach, C.C., Graber, J.W., Niswender, G.D. and Naibandov, A.V. (1968b). Luteotropic properties of some pituitary hormones in nonpregnant and pregnant hypophysectomized ewes. *Endocrinology* 82:818-824.
- Kasson, B.G. and Hsueh, A.J.W. (1985a). Cholinergic inhibition of follicle-stimulating hormone-induced progesterone production by cultured rat granulosa cells. *Biol. Reprod.* 33:1158-1167.
- Kasson, B.G. and Hsueh, A.J.W. (1985b). Nicotinic cholinergic agonists inhibit androgen biosynthesis by cultured rat testicular cells. *Endocrinology* 117:1874-1880.
- Kawai, Y. and Clark, M.R. (1985). Phorbol ester regulation of rat granulosa cell prostaglandin and progesterone accumulation. *Endocrinology* 116:2320-2326.
- Kawakami, M., Kubo, K., Uemura, T., Nagase, M. and Hayashi, R. (1981). Involvement of ovarian innervation in steroid secretion. *Endocrinology* 109:136-145.
- Keyes, P.L. and Naibandov, A.V. (1967). Maintenance and function of corpora lutea in rabbits depends on estrogen. *Endocrinology* 80:938-943.
- Khan-Dawood, F.S. and Dawood, M.Y. (1983). Human ovaries contain immunoreactive oxytocin. *J. Clin. Endocrinol. Metab.* 57:1129-1132.
- Khan, M.I. and Gibori, G. (1984). Catechol estrogens and their role in luteal steroidogenesis. *Biol. Reprod.* 30 Suppl. 1:194 (abstract).

- Kimball, F.A. and Hansel, W. (1974). Estrogen cytosol binding proteins in bovine endometrium and corpus luteum. *Biol. Reprod.* 11:566-577.
- Kimball, F.A. and Lauderdale, J.W. (1975). Prostaglandin E_1 and F_2^{α} specific binding in bovine corpora lutea: comparison with luteolytic effects. *Prostaglandins* 10:313-331.
- Knecht, M. and Catt, K.J. (1981). Gonadotropin-releasing hormone: regulation of adenosine 3',5'-monophosphate in ovarian granulosa cells. *Science* 214:1346-1348.
- Knecht, M., Katz, M.S. and Catt, K.J. (1981). Gonadotropin-releasing hormone inhibits cyclic nucleotide accumulation in cultured rat granulosa cells. *J. Biol. Chem.* 256:34-36.
- Knecht, M. and Catt, K.J. (1983). Modulation of cAMP-mediated differentiation in ovarian granulosa cells by epidermal growth factor and platelet derived growth factor. *J. Biol. Chem.* 258:2789-2794.
- Koos, R.D. and Hansel, W. (1981). The large and small cells of the bovine corpus luteum: Ultrastructural and functional differences. In: *Dynamics of Ovarian Function*. (N.B. Schwartz and M. Hunzicker-Dunn, eds.). Raven Press, New York, pp. 197-203.
- Koos, R.D. and Clark, M.R. (1982). Production of 6-keto-prostaglandin F_1^{α} by rat granulosa cells *in vitro*. *Endocrinology* 111:1513-1518.
- Kovanen, P.T., Goldstein, J.L. and Brown, M.S. (1978). High levels of 3-hydroxy-3-methylglutaryl coenzyme A reductase activity and cholesterol synthesis in the ovary of the pregnant rabbit. *J. Biol. Chem.* 253:5126-5132.
- Krebs, E.G. and Beavo, J.A. (1979). Phosphorylation-dephosphorylation of enzymes. *Ann. Rev. Biochem.* 48:923-959.
- Kumar, A. (1984). Absence of effect of a gonadotropin releasing hormone agonist on progesterone production in bovine luteal cells *in vitro*. *IRSC Med. Sci.* 12:114.
- Kuo, J.F., Schatzman, R.C., Turner, R.S. and Mazzei, G.J. (1984). Phospholipid-sensitive Ca^{2+} -dependent protein kinase: a major protein phosphorylation system. *Molec. Cellular Endocrinol.* 35:65-73.
- Ladenheim, R.G., Tesone, M. and Charreau, E.H. (1984). Insulin action and characterization of insulin receptors in rat luteal cells. *Endocrinology* 115:752-756.
- Larsson, L.I., Fahrenkrug, J. and Schaffalitzky de Muckadell, O.B. (1977). Vasoactive intestinal peptide occurs in nerves of the female genitourinary tract. *Science* 197:1374-1375.
- Laszlovsky, I. and Erdo, S.L. (1983). Characterization of β -adrenergic

- receptors in rat ovary. *Eur. J. Pharmacol.* 96:101-104.
- Lawrence, I.E. and Burden, H.W. (1980). The origin of the extrinsic adrenergic innervation to the rat ovary. *Anat. Rec.* 196:51-62.
- Lefkowitz, R.J. (1976). β -adrenergic receptors: recognition and regulation. *N. Engl. J. Med.* 295:323-328.
- Legrand, C. and Maltier, J.P. (1986). Evidence for a noradrenergic transmission in the control of parturition in the rat. *J. Reprod. Fert.* 76:415-424.
- Lemon, M. and Loir, M. (1977). Steroid release in vitro by two luteal cell types in the corpus luteum of the pregnant sow. *J. Endocrinol.* 72:351-359.
- LePere, R.H., Benoit, P.E., Hardy, R.C. and Goldzieher, J.W. (1966). The origin and function of the ovarian nerve supply in the baboon. *Fert. Steril.* 17:68-75.
- Levine, J., Lloyd, C.W., Lobotsky, J. and Friedrick, E.H. (1967). The effect of epinephrine on testosterone production. *Acta Endocrinol. Copenh.* 55:184-191.
- Lincoln, G.A. (1984). The pineal gland. In: *Reproduction in Mammals, Book 3: Hormonal Control of Reproduction*. (C.R. Austin and R.V. Short, eds.). Cambridge University Press, London, pp.52-75.
- Lim, A.T., Lolait, S., Barlow, J.W., Wai Sum, O., Zols, I., Toh, B.H. and Funder, J.W. (1983). Immunoreactive β -endorphin in sheep ovary. *Nature* 303:709-711.
- Ling, W.Y. and Marsh, J.M. (1977). Reevaluation of the role of cyclic adenosine 3',5'-monophosphate and protein kinase in the stimulation of steroidogenesis by luteinizing hormone in bovine corpus luteum slices. *Endocrinology* 100:1571-1578.
- Lloyd, T. and Welsz, J. (1978). Direct inhibition of tyrosine hydroxylase activity by catechol estrogens. *J. Biol. Chem.* 253:4841-4843.
- Loeb, L. (1917). The relations of the ovary to the uterus and mammary gland from the experimental aspect. *Surg. Gynecol. Obstet.* 25:300.
- Lolait, S.J., Autelitano, D.J., Lim, A.T.W., Smith, A.I., Toh, B.H. and Funder, J.W. (1985). Ovarian immunoreactive β -endorphin and estrous cycle in the rat. *Endocrinology* 117:161-168.
- Louvet, J.P., Balslic, M., Bayard, F. and Boulard, C. (1977). Glucocorticoid receptors in rat ovarian granulosa cell cytosol. *Endocrinology* 100, Suppl. 1:601 (abstract).
- MacLusky, N.J., Barnea, E.R., Clark, C.R. and Naftolin, F. (1983). Catechol estrogens and estrogen receptors. In: *Catechol Estrogens*. (G.R. Merriam and M.B. Lipsett, eds.). Raven Press, New York,

pp.151-170.

- MacPhee, A.A., Cole, F.E. and Rice, B.F. (1975). The effect of melatonin on steroidogenesis by the human ovary in vitro. J. Clin. Endocrinol. Metab. 40:688-696.
- Mahayan, D.K. and Little, A.B. (1978). Specific cortisol binding protein in porcine follicular fluid. Biol. Reprod. 17:834-842.
- Makris, A., Yasumizu, T., Elkind-Hirsch, K., Carraway, R., Leeman, S.E. and Ryan, R.J. (1982). Substance P and neurotensin in the ovary and pituitary of the cyclic golden hamster. Biol. Reprod. 26 Suppl. 1:123 (abstract).
- Maltier, J.P., Legrand, C. and Corazza, S. (1984). Etude des recepteurs α -adrenergiques dans le myometre de rate preparturiente. Path. Biol. 32:878-883.
- Marsh, J.M. (1970). The stimulatory effect of luteinizing hormone on adenylyl cyclase in the bovine corpus luteum. J. Biol. Chem. 245:1596-1603.
- Marsh, J.M. (1971). The effect of prostaglandins on the adenylyl cyclase of the bovine corpus luteum. Ann. NY Acad. Sci. 180:416-425.
- Marsh, J.M. (1976). The role of cyclic AMP in gonadal steroidogenesis. Biol. Reprod. 14:30-53.
- Marsh, J.M., Butcher, R.W., Savard, K. and Sutherland, E.W. (1966). The stimulatory effect of luteinizing hormone on adenosine 3',5'-monophosphate accumulation in corpus luteum slices. J. Biol. Chem. 241:5436-5440.
- Marsh, J.M. and Savard, K. (1966). Studies on the mode of action of luteinizing hormone on steroidogenesis in the corpus luteum in vitro. J. Reprod. Fert. Suppl. 1:113-116.
- Marshall, J.M. (1970). Adrenergic innervation of the female reproductive tract: anatomy, physiology and pharmacology. Rev. Physiol. Biochem. Pharmacol. 62:5-67.
- Mason, N.R., Marsh, J.M. and Savard, K. (1962). An action of gonadotropin in vitro. J. Biol. Chem. 237:1801-1805.
- May, J.V., McCarty, K., Reichert, L.E. and Schomberg, D.W. (1980). Follicle-stimulating hormone-mediated induction of functional luteinizing hormone/human chorionic gonadotropin receptors during monolayer culture of porcine granulosa cells. Endocrinology 107:1041-1049.
- May, J.V. and Schomberg, D.W. (1981). Granulosa cell differentiation in vitro: effect of insulin on growth and functional integrity. Biol. Reprod. 25:421-431.
- McCracken, J.A., Carlson, J.C., Glew, M.E., Goding, J.R. and Baird, D.T.

- (1972). Prostaglandin $F_{2\alpha}$ identified as a luteolytic hormone in sheep. *Nature New Biol.* 238:129-134.
- McCracken, J.A. (1980). Hormone receptor control of prostaglandin $F_{2\alpha}$ secretion by ovine uterus. In: *Advances in Prostaglandin and Thromboxane Research.* (b. Samuelsson, P.W. Ramwell and R. Paoletti, eds.). Raven Press, New York, pp.1329-1344.
- McNutt, G.W. (1924). The corpus luteum of the ox ovary in relation to the oestrous cycle. *J. Am. Vet. Med. Assoc.* 65:556-561.
- Menon, K.M.J. (1973). Purification and properties of a protein kinase from bovine corpus luteum that is stimulated by cyclic adenosine 3',5'-monophosphate and luteinizing hormone. *J. Biol. Chem.* 248:494-501.
- Michell, R.H. (1975). Inositol phospholipids and cell surface receptor function. *Biochim. Biophys. Acta* 415:81-147.
- Milvae, R.A. and Hansel, W. (1980). The effects of prostacyclin and 6-keto-PGF $_{1\alpha}$ on bovine plasma progesterone and LH concentrations. *Prostaglandins* 20:641-647.
- Milvae, R.A., Allia, H.W. and Hansel, W. (1983). Methylation in bovine luteal cells as a regulator of luteinizing hormone action. *Biol. Reprod.* 29:849-855.
- Milvae, R.A. and Hansel, W. (1983). Prostacyclin, prostaglandin $F_{2\alpha}$ and progesterone production by bovine luteal cells during the estrous cycle. *Biol. Reprod.* 29:1063-1068.
- Milvae, R.A. and Hansel, W. (1985). Inhibition of bovine luteal function by indomethacin. *J. Anim. Sci.* 60:528-531.
- Mitchell, G.A.G. (1938). The innervation of the ovary, uterine tube and epididymis. *J. Anat.* 72:508-515.
- Moger, W.H. and Anakive, O.O. (1986). Propranolol inhibits the compensatory increase in androgen secretion after unilateral orchidectomy in rats. *J. Reprod. Fert.* 76:251-256.
- Mori, M. and Marsh, J.M. (1982). The site of luteinizing hormone stimulation of steroidogenesis in mitochondria of the rat corpus luteum. *J. Biol. Chem.* 257:6178-6183.
- Mossman, H.W. and Duke, K.L. (1973). Some comparative aspects of the mammalian ovary. In: *Handbook of Physiology, Section 7: Endocrinology Vol. 1, Part 2.* (S.R. Geiger, ed.). Physiological Society, Washington, D.C., pp.389-402.
- Mukhopadhyay, A.K., Kumar, A., Tams, R., Bohnet, H.G. and Leidenberger, F.A. (1984). Oxytocin and vasopressin have no effect on progesterone production and cyclic AMP accumulation by rat luteal cells *in vitro*. *J. Reprod. Fert.* 72:137-141.

- Naumenko, E.V. (1985). Role of catecholamines and serotonin in the control of the hypothalamic-pituitary-testicular complex- A review. *Biogenic Amines* 3:173-190.
- Nett, T.M., McClellan, M.C. and Niswender, G.D. (1976). Effects of prostaglandins on the ovine corpus luteum: Blood flow, secretion of progesterone and morphology. *Biol. Reprod.* 15:66-78.
- Niedel, J.E., Kuhn, L.J. and Vandenbark, G.R. (1983). Phorbol diester receptor copurifies with protein kinase C. *Proc. Natl. Acad. Sci.* 80:36-40.
- Niilius, S.J., Bergquist, C. and Wide, L. (1978). Inhibition of ovulation in women by chronic treatment with a stimulatory LRH analogue, a new approach to birth control? *Contraception* 17:537-542.
- Nishizuka, Y. (1984). Turnover of inositol phospholipids and signal transduction. *Science* 225:1365-1370.
- Niswender, G.D., Schwall, R.H., Fitz, T.A., Farin, C. and Sawyer, H.R. (1985). Regulation of ovarian function in domestic ruminants: New concepts. *Recent Prog. Horm. Res.* 41:101-152.
- Norjavaara, E., Selstam, G. and Ahren, K. (1982). Catecholamine stimulation of cyclic AMP and progesterone production in rat corpora lutea of different ages. *Acta Endocrinol. Copenh.* 100:613-622.
- Norjavaara, E., Sestam, G., Damberg, J.E. and Johansson, B.M. (1983). In vivo effect of noradrenaline on the cyclic AMP level in rat corpora lutea. *Acta Physiol. Scand.* 119:113-116.
- Norjavaara, E., Rosberg, S., Gafvels, M. and Selstam, G. (1984). β -adrenergic receptor concentration in corpora lutea of different ages obtained from pregnant mare serum gonadotropin-treated rats. *Endocrinology*. 114:2154-2159.
- O'Byrne, E.M., Filitcraft, J.F., Sawyer, W.K., Hochman, J., Weiss, G., and Steinmetz, B.G. (1978). Relaxin bioactivity and immunoactivity in human corpora lutea. *Endocrinology* 102:1641-1644.
- Ojeda, S.R., Costa, M.E., Kalz, K.H. and Hersh, L.B. (1985). Evidence for the existence of substance P in the prepubertal rat ovary. I. Biochemical and physiological studies. *Biol. Reprod.* 33:286-295.
- O'Shea, J.D., Cran, D.G. and Hay, M.F. (1979). The small luteal cell of the sheep. *J. Anat.* 128:239-251.
- Owman, Ch., Rosengren, E. and Sjöberg, N.O. (1967). Adrenergic innervation of the human female reproductive organs: A histochemical and chemical investigation. *Obstet. Gynecol.* 30:763-773.
- Owman, C., Sjöberg, N.O., Svensson, K.G. and Wallies, B. (1975). Autonomic nerves mediating contractility in the human graafian follicle. *J. Reprod. Fert.* 45:553-556.

- Papka, R.E., Cotton, J.P. and Taurig, H.H. (1985). Comparative distribution of neuropeptide tyrosine, vasoactive intestinal polypeptide, substance P-immunoreactive, acetylcholinesterase-positive and noradrenergic nerves in the reproductive tract of the female rat. *Cell Tissue Res.* 242:475-490.
- Pate, J.P. and Condon, W.A. (1984). Effects of prostaglandin F_2^α on agonist-induced progesterone production in cultured bovine luteal cells. *Biol. Reprod.* 31:427-435.
- Peat, F. and Kinson, G.A. (1971). Testicular steroidogenesis *in vitro* in the rat in response to blinding, pinealectomy and to the addition of melatonin. *Steroids* 17:251-264.
- Perkins, S.N., Cronin, M.J. and Veldhuis, J.D. (1986). Properties of β -adrenergic receptors on porcine corpora lutea and granulosa cells. *Endocrinology* 118:998-1005.
- Pieper, D.R., Richards, J.S. and Marshall, J.C. (1981). Ovarian gonadotropin-releasing hormone (GnRH) receptors: characterization, distribution and induction by GnRH. *Endocrinology* 108:1148-1155.
- Pitzel, L., Welp, K., Holtz, W. and König, A. (1984). Neurohypophyseal hormones in the corpus luteum of the pig. *Neuroendocrinol. Lett.* 6:1-5.
- Poff, J.P., Fairchild, D.F. and Condon, W.A. (1987). Effects of antibiotics and medium supplements on steroidogenesis in cultured bovine luteal cells. *Biol. Reprod.*, submitted.
- Polan, M.L., Decherney, A.H., Haseltine, F.D., Meyer, H.C. and Behrman, H.R. (1983). Adenosine amplifies FSH action in granulosa cells and LH action in luteal cells of the rat and human ovary. *J. Clin. Endocrinol.* 56:288-294.
- Powell, W.S., Hammarstrom, S. and Samuelsson, B. (1974a). Prostaglandin F_2^α receptor in ovine corpora lutea. *Eur. J. Biochem.* 41:103-107.
- Powell, W.S., Hammarstrom, S., Samuelsson, B. and Sjöberg, B. (1974b). Prostaglandin F_2^α receptor in human corpora lutea. *Lancet* 1:1120.
- Priedkalns, J. and Weber, A.F. (1968). Ultrastructural studies of the bovine Graafian follicle and corpus luteum. *Zeit. Zell. Mik. Anat.* 91:554-573.
- Quagliarello, J., Goldsmith, L., Steinetz, B., Lustig, D.S. and Weis, G. (1980). Induction of relaxin secretion in nonpregnant women by human chorionic gonadotropin. *J. Clin. Endocrinol. Metab.* 51:74-77.
- Quirk, S.J., Willcox, D.L., Parry, D.M. and Thornburn, G.D. (1979). Subculture location of progesterone in the bovine corpus luteum: a biochemical, morphological and cytochemical investigation. *Biol. Reprod.* 20:1133-1145.

- Ranta, T., Knecht, M., Kody, M. and Catt, K.J. (1982). Gonadotropin releasing hormone receptors in cultured rat granulosa cells: Mediation of the inhibitory and stimulatory actions of gonadotropin releasing hormone. *Molec. Cellular Endocrinol.* 27:233-240.
- Rao, Ch.V. (1974). Characterization of prostaglandin receptors in the bovine corpus luteum cell membranes. *J. Biol. Chem.* 249:7203-7209.
- Rao, Ch.V. (1975). The presence of discrete receptors for prostaglandin $F_{2\alpha}$ in the cell membranes of bovine corpora lutea. *Biochem. Biophys. Res. Comm.* 64:416-424.
- Rao, Ch.V., Fields, M.J., Chen, T.T., Abel, J.H. and Edgerton, L.A. (1983). Change in gonadotropin-binding sites in intracellular organelles and plasma membranes during luteal growth, development and regression. *Exp. Cell Res.* 144:285-295.
- Ratner, A., Sanborn, C.R. and Weiss, G.K. (1980a). β -Adrenergic stimulation of cAMP and progesterone in rat ovarian tissue. *Am. J. Physiol.* 239:E139-E143.
- Ratner, A., Weiss, G.K. and Sanborn, C.R. (1980b). Stimulation by β -adrenergic receptors of the production of cyclic AMP and progesterone in rat ovarian tissue. *J. Endocrinol.* 87:123-129.
- Reiter, R.J. (1984). Pineal indoles: production, secretion and actions. In: *Neuroendocrine Perspectives, Volume 3.* (E.E. Muller and R.M. MacLeod, eds.). Elsevier Science Publishers, New York, pp.345-377.
- Rice, G.E., Jenkin, G. and Thorburn, G.D. (1986). Comparison of particle-associated progesterone and oxytocin in the ovine corpus luteum. *J. Endocrinol.* 108:109-116.
- Richards, J.S. (1974). Estradiol binding to rat corpora lutea during pregnancy. *Endocrinology* 95:1046-1053.
- Richardson, M.C. and Masson, G.M. (1980). Progesterone production by dispersed cells from human corpus luteum: stimulation by gonadotrophins and prostaglandin $F_{2\alpha}$ lack of response to adrenaline and isoprenaline. *J. Endocrinol.* 87:247-254.
- Richardson, M.C. and Masson, G.M. (1985). Lack of direct inhibitory effect of oxytocin on progesterone production by dispersed cells from human corpus luteum. *J. Endocrinol.* 104:149-151.
- Rivier, G., Rivier, J. and Vale, W. (1978). Chronic effects of [D-Trp⁶-Pro⁹-NEt] luteinizing hormone-releasing factor on reproductive processes in the female rat. *Endocrinology* 103:2299-2305.
- Rodbell, M. (1980). The role of hormone receptors and GTP-regulatory proteins in membrane transduction. *Nature* 284:17-22.
- Rodgers, R.J., O'Shea, J.D. and Findlay, J.K. (1983). Progesterone production *in vitro* by small and large ovine luteal cells. *J. Reprod. Fert.* 69:113-124.

- Rodgers, R.J., O'Shea, J.D. and Bruce, N.W. (1984). Morphometric analysis of the cellular composition of the ovine corpus luteum. *J. Anat.* 138:757-769.
- Rommerts, F.F.G., Molenaar, R., Hoogerbrugge, J.W. and Van Der Molen, H.J. (1984). Development of adenosine responsiveness after isolation of leydig cells. *Biol. Reprod.* 30:842-847.
- Saez, J.H., Morera, A.M., Haour, F. and Evain, D. (1977). Effect of in vivo administration of dexamethasone, corticotropin and human chorionic gonadotropin on steroidogenesis and protein and DNA synthesis of testicular interstitial cells in prepuberal rats. *Endocrinology* 101:1256-1263.
- Savard, K., Marsh, J.M. and Rice, B.F. (1965). Gonadotropins and ovarian steroidogenesis. *Recent Prog. Horm. Res.* 21:285-365.
- Savard, K. (1973). The biochemistry of the corpus luteum. *Biol. Reprod.* 8:183-202.
- Savion, N. and Gospodarowicz, D. (1980). Patterns of cellular peptide synthesis by cultured bovine granulosa cells. *Endocrinology* 107:1798-1807.
- Sawyer, H.R., Abel, J.H. Jr., McClellan, M.C., Schmitz, M. and Niswender, G.D. (1979). Secretory granules and progesterone secretion by ovine corpora lutea in vitro. *Endocrinology* 104:476-486.
- Schofield, B.M. (1952). The innervation of the cervix and cornu uteri in the rabbit. *J. Physiol.* 117:317-324.
- Schuler, L.A., Toaff, M.A. and Strauss, J.F. III (1981). Regulation of ovarian cholesterol metabolism: control of 3-hydroxy-3-methylglutaryl coenzyme A reductase and acyl coenzyme A: cholesterol acyltransferase. *Endocrinology* 108:1476-1486.
- Seguin, B.E., Oxender, W.D. and Britt, J.H. (1977). Effect of human chorionic gonadotropin and gonadotropin-releasing hormone on corpus luteum function and estrous cycle duration in dairy heifers. *Am. J. Vet. Res.* 38:1153-1156.
- Selfart, K.H. and Hansel, W. (1968). Some characteristics and optimum incubation conditions of in vitro progesterone synthesis by bovine corpora lutea. *Endocrinology* 82:232-240.
- Shaeffer, J.M. and Hsueh, A.J.W. (1982). Identification of gamma-aminobutyric acid and its binding sites in the rat ovary. *Life Sci.* 30:1599-1604.
- Shaha, C., Margioris, A., Liotta, A.S., Krieger, D.T. and Bardin, C.W. (1984). Demonstration of immunoreactive δ -endorphin and γ_3 -melanocyte stimulating hormone related peptides in the ovaries of neonatal, cyclic and pregnant mice. *Endocrinology* 115:378-384.

- Shinohara, C., Knecht, M., Feng, P. and Catt, K.J. (1986). Activation of protein kinase C potentiates cyclic AMP production and stimulates steroidogenesis in differentiated ovarian granulosa cells. *J. Steroid Biochem.* 24:161-168.
- Shemesh, M. and Hansel, W. (1975a). Stimulation of prostaglandin synthesis in bovine ovarian tissue by arachidonic acid and luteinizing hormone. *Biol. Reprod.* 13:448-452.
- Shemesh, M. and Hansel, W. (1975b). Arachidonic acid and bovine corpus luteum function. *Proc. Exp. Biol. Med.* 148:123-126.
- Sherwood, O.D. and O'Byrne, E.M. (1974). Purification and characterization of porcine relaxin. *Arch. Biochem. Biophys.* 160:185-196.
- Sherwood, O.D. and Rutherford, J.E. (1981). Relaxin immunoreactivity levels in ovarian extracts obtained from rats during various reproductive states and from adult cycling pigs. *Endocrinology* 108:1171-1177.
- Sherwood, O.D. and Downing, S.J. (1983). The chemistry and physiology of relaxin. In: *Factor Regulating Ovarian Function*. (G.S. Greenwald and P.F. Terranova, eds.). Raven Press, New York, pp.381-410.
- Short, R.V. (1964). Ovarian steroid synthesis and secretion in vivo. *Rec. Prog. Horm. Res.* 20:303-332.
- Simmons, K.R. and Hansel, W. (1964). Nature of the luteotropic hormone in the bovine. *J. Anim. Sci.* 23:136-141.
- Simpson, E.R., McCarthy, J.L. and Peterson, J.A. (1978). Evidence that the cycloheximide-sensitive site of adrenocorticotrophic hormone action is in the mitochondria. Changes in pregnenolone formation, cholesterol content, and electron paramagnetic resonance spectra of cytochrome P-450. *J. Biol. Chem.* 253:3135-3139.
- Sinha, A.A., Seal, U.S. and Doe, R.P. (1971). Ultrastructure of the corpus luteum of the white-tailed deer during pregnancy. *Am. J. Anat.* 132:189-206.
- Sjoberg, N.O. (1967). The adrenergic transmitter of the female reproductive tract: distribution and function changes. *Acta Physiol. Scand. Suppl.* 305:5-32.
- Sulimovici, S.I. and Boyd, E.S. (1969). The cholesterol side chain cleavage enzymes in steroid hormone-producing tissues. *Vitam. Horm.* 27:199-234.
- Snook, R.B., Brunner, M.A., Saatman, R.R. and Hansel, W. (1969). The effect of antisera to bovine LH in hysterectomized and intact heifers. *Biol. Reprod.* 1:49-58.
- Spielvogel, A.M., Mednicks, M.I., Eppenberger, U. and Jungmann, R.A.

- (1977). Evidence for the identity of nuclear and cytoplasmic adenosine 3',5'-monophosphate-dependent protein kinase from porcine ovaries and nuclear translocation of the cytoplasmic enzyme. *Eur. J. Biochem.* 73:199-212.
- Stefenson, A., Owman, Ch., Sjöberg, N.O., Spörng, B. and Wallis, B. (1981). Comparative study of the autonomic innervation of the mammalian ovary, with particular regard to the follicular system. *Cell Tissue Res.* 215:47-62.
- Sternquist, M., Emson, P., Owman, Ch., Sjöberg, N.O., Sundler, F. and Tatemoto, K. (1983). Neuropeptide Y in the female reproductive tract of the rat. Distribution of nerve fibers and motor effects. *Neurosci. Lett.* 9:279-284.
- Stouffer, R.L., Tyrey, L. and Schomberg, D.W. (1976). Changes in [¹²⁵I]-labeled human chorionic gonadotropin (hCG) binding to porcine granulosa cells during follicular development and cell culture. *Endocrinology* 99:516-525.
- Strauss, J.F. III, Schuler, L.A., Rosenblum, M.F. and Tanaka, T. (1981). Cholesterol metabolism of ovarian tissue. *Adv. Lipid Res.* 18:99-157.
- Tamaoki, B.I. (1973). Steroidogenesis and cell structure. Biochemical pursuit of sites of steroid biosynthesis. *J. Steroid Biochem.* 4:89-118.
- Tan, G.J.S., Tweedale, R. and Biggs, J.S.G. (1982a). Effects of oxytocin on the bovine corpus luteum of pregnancy. *J. Reprod. Fert.* 66:75-78.
- Tan, G.J.S., Tweedale, R. and Biggs, J.S.G. (1982b). Oxytocin may play a role in the control of the human corpus luteum. *J. Endocrinol.* 95:65-70.
- Tan, G.J.S. and Biggs, J.S.G. (1983). Absence of effect of LH-RH on progesterone production by human luteal cells *in vitro*. *J. Reprod. Fert.* 67:411-413.
- Tesone, M., Ladenheim, R.G., Oliveira-Filho, R.M., Chlauzzi, V.A., Foglia, V.G. and Charreau, E.H. (1983). Ovarian dysfunction in streptozotocin-induced diabetic rats. *Proc. Soc. Exp. Biol. Med.* 174:123-127.
- Traurig, H., Papka, R.E. and Cotton, J.P. (1984). Neuropeptide Y (NPT)-like immunoreactivity in the female reproductive system. *Neurosci.* 10:604 (abstract).
- Trzeciak, W.H. and Boyd, G.S. (1973). The effect of stress induced by ether anesthesia on cholesterol content and cholesteryl-esterase activity in rat-adrenal cortex. *Eur. J. Biochem.* 37:327-333.
- Tsong, S.D., Phillips, D., Halmi, N., Liotta, A.S., Margioris, A., Bardin, C.W. and Krieger, D.T. (1982a). ACTH and β -endorphin-

- related peptides are present in multiple sites in the reproductive tract of the male rat. *Endocrinology* 110:2204-2206.
- Tsong, S.D., Phillips, D., Halmi, N., Krieger, D.T. and Bardin, C.W. (1982b). δ -endorphin is present in the male reproductive tract of five species. *Biol. Reprod.* 27:755-764.
- Unsicker, K. (1974). Qualitative and quantitative studies on the innervation of the corpus luteum of the rat and pig. *Cell Tissue Res.* 152:513-524.
- Ursely, J. and Leymarie, P. (1979). Varying response to luteinizing hormone of two luteal cell types isolated from bovine corpus luteum. *J. Endocrinol.* 83:303-310.
- Vaitukaitis, J.L., Lee, C.Y., Ebersole, E.R. and Lerario, A.C. (1975). New evidence for an acute role of protein kinase in hCG action. *Endocrinology* 97:215-222.
- Van Orden, D.E., Goodale, D.B., Baker, H.A., Farley, D.B. and Bhatnagar, R.K. (1980). Uterine catecholamines and prostaglandins during the estrous cycle of the rat. *Endocrinology* 106:1650-1654.
- Veldhuis, J.D., Kolp, L.A., Toaff, M.E., Strauss, J.F. and Demers, L.M. (1983). Mechanisms subserving the trophic actions of insulin on ovarian cells: in vitro studies using swine granulosa cells. *J. Clin. Invest.* 72:1046-1057.
- Veldhuis, J.D. and Furlanetto, R.W. (1985). Trophic actions of human somatomedin C/insulin-like growth factor I on ovarian cells: in vitro studies with swine granulosa cells. *Endocrinology* 116:1235-1242.
- Veldhuis, J.D., Rodgers, R.J., Dee, A. and Simpson, E.R. (1986). The insulin-like growth factor, somatomedin-C, induces the synthesis of cholesterol side-chain cleavage cytochrome P-450 and adrenodoxin in ovarian cells. *J. Biol. Chem.* 261:2499-2502.
- Verdesca, A.S., Westermann, C.D., Crampton, R.S., Black, W.C., Nedeljkovic, R.I. and Hilton, J.C. (1961). Direct adrenocortical stimulatory effect of serotonin. *Am. J. Physiol.* 210:1065-1067.
- Virutamasen, P., Wright, K.H. and Wallach, E.E. (1973). Monkey ovarian contractility-the relationship to ovulation. *Fert. Steril.* 24:763-769.
- Virutamsen, P., Smithasiri, Y. and Fuch, A.R. (1976). Intra-ovarian pressure changes during ovulation in rabbits. *Fert. Steril.* 27:188-192.
- Vlodarsky, I., Brown, K.D. and Gospodarowicz, D. (1978). A comparison of the binding of epidermal growth factor to cultured granulosa and luteal cells. *J. Biol. Chem.* 253:3744-3750.
- Vogt, M. (1943). The output of cortical hormone by the mammalian suprarenal. *J. Physiol.* 102:341-356.

- Wallis, B., Edvinsson, L., Owman, Ch., Sjöberg, N.O. and Svensson, K.G. (1975). Evidence for a neuromuscular mechanism involved in the contractility of the ovarian follicular wall: fluorescence and electron microscopy and effects of tyramine on follicular strips. *Biol. Reprod.* 12:239-248.
- Wallis, B., Edvinsson, L., Owman, Ch., Sjöberg, N.O. and Spörng, B. (1976). Cholinergic nerves and receptors mediating contraction of the Graafian follicle. *Biol. Reprod.* 15:565-572.
- Wallis, B., Groschel-Stewart, V., Owman, Ch., Sjöberg, N.O. and Unsicker, K. (1978). Fluorescence histochemical demonstration of a relationship between adrenergic nerves and cells containing actin and myosin in the rat ovary, with special regard to the follicular wall. *J. Reprod. Fert.* 52:175-178.
- Walsh, D.A. and Ashly, C.D. (1973). Protein kinases: Aspects of their regulation and diversity. *Rec. Prog. Horm. Res.* 29:329-353.
- Warbritton, V. (1934). The cytology of the corpora lutea of the ewe. *J. Morph.* 56:181-202.
- Wathes, D.C. and Swann, R.W. (1982). Is oxytocin an ovarian hormone? *Nature* 297:225-227.
- Wathes, D.C., Swann, R.W., Pickering, B.T., Porter, D.G., Hall, M.G.R. and Drife, J.O. (1982). Neurohypophyseal hormones in the human ovary. *Lancet* 2:410-412.
- Wathes, D.C., Swann, R.W., Birkett, S.D., Porter, D.G., and Pickering, B.T. (1983). Characterization of oxytocin, vasopressin and neurophysin from the bovine corpus luteum. *Endocrinology* 113:693-698.
- Watkins, W.B. (1983). Immunohistochemical localization of neurophysin and oxytocin in the sheep corpora lutea. *Neuropeptides* 4:51-54.
- Weiss, G.K., Dail, W.G. and Ratner, A. (1982). Evidence for a direct neural control of ovarian steroidogenesis in rats. *J. Reprod. Fert.* 65:507-511.
- Welsh, T.H. and Hsueh, A.J.W. (1982). Mechanism of the inhibitory action of epidermal growth factor on testicular androgen biosynthesis in vitro. *Endocrinology* 110:1498-1506.
- Wilkinson, R.F., Anderson, E. and Aalberg, J. (1976). Cytological observations of dissociated rat corpus luteum. *J. Ultrastruct. Res.* 57:168-184.
- Willcox, D.L. (1983). Two binding proteins for progesterone in the bovine corpus luteum. *Biol. reprod.* 29:487-497.
- Williams, M.T. and Marsh, J.M. (1978). Estradiol inhibition of luteinizing hormone-stimulated progesterone synthesis in isolated

bovine luteal cells. *Endocrinology* 103:1611-1618.

- Williams, M.T., Clark, M.R., Ling, W.Y., LeMarle, W.J., Caron, M.G. and Marsh, J.M. (1978). Role of cyclic AMP in the action of luteinizing hormone on steroidogenesis in the corpus luteum. In: *Advances in Cyclic Nucleotide Research*. (W.J. George and L.J. Ignarro, eds.). Raven Press, New York, pp.573-582.
- Willingham, M.C. and Pastan, I. (1984). Endocytosis and membrane traffic in cultured cells. *Rec. Prog. Horm. Res.* 40:569-587.
- Willbank, J.N., Rothlisberger, J.A. and Zimmerman, D.R. (1961). Effect of human chorionic gonadotropin on maintenance of the corpus luteum and embryonic survival in the cow. *J. Anim. Sci.* 20:827-829.
- Wolff, J. and Hope-Cook, G. (1977). Activation of steroidogenesis and adenylate cyclase by adenosine and Leydig tumor cells. *J. Biol. Chem.* 252:687-693.
- Wurtman, R.J., Chu, F.W. and Axelrod, J. (1963). Relation between the oestrous cycle and the binding of catecholamines in the rat uterus. *Nature* 198:547-549.
- Wurtman, R.J., Axelrod, J. and Potter, I.T. (1964). The uptake of 3H -melatonin in endocrine and nervous tissues and the effects of constant light exposure. *J. Pharmacol. exp. Ther.* 143:314-321.
- Zimniski, S.J., Rorke, E.A., Helmy, R.S. and Vaitukaitis, J.L. (1982). Characterization of human chorionic gonadotropin internalization by pseudopregnant rat ovary. *Endocrinology* 111:635-640.
- Zor, U., Strulovici, B. and Lindner, H.R. (1978). Implication of microtubules and microfilaments in the response of the ovarian adenylate cyclase-cyclic AMP system to gonadotropin and prostaglandin E_2 . *Biochem. Biophys. Res. Comm.* 80:983-992.
- Zsolnai, B., Varga, B. and Horvath, E. (1982). Increase of ovarian progesterone secretion by β_2 -adrenergic stimulation in oestrous rats. *Acta Endocrinol. Copenh.* 101:268-272.

EXPERIMENT ONE

Serotonin-induced stimulation of progesterone
production by cow luteal cells in vitro

P. J. Battista and W. A. Condon

Department of Animal and Nutritional Sciences
The University of New Hampshire
Durham, New Hampshire 03824

Published in the Journal of Reproduction
and Fertility 76:231-238 (1986).

Summary

The addition of acetylcholine or histamine (10^{-7} to 10^{-4} M), γ -aminobutyric acid, a dopamine agonist, and melatonin (10^{-7} to 10^{-5} M) did not alter basal or LH-stimulated progesterone production ($P>0.05$). The addition of the specific β_2 -adrenergic agonists terbutaline and salbutamol did not significantly elevate progesterone production. Treatment of luteal cells with serotonin (5-HT), 10^{-6} to 10^{-4} M increased the production of progesterone ($P<0.05$). This stimulated production was inhibited by the addition of mianserin (10^{-5} M, a 5-HT antagonist; $P<0.05$). Isoproterenol (10^{-7} to 10^{-4} M) also resulted in significant increases in progesterone production ($P<0.05$). The combined treatments of 5-HT + LH, Isoproterenol + LH, or Isoproterenol + 5-HT did not result in a further increase in progesterone above that observed in response to LH or Isoproterenol alone ($P>0.05$). The Isoproterenol-induced progesterone production could not be blocked by butoxamine (10^{-5} M, a β_2 -antagonist), or practolol (10^{-5} M, a β_1 -antagonist), but was inhibited by propranolol (10^{-5} M, a general β -antagonist; $P<0.05$). The response to Isoproterenol was unaffected by mianserin (10^{-5} M). These results demonstrate a possible role for 5-HT in the stimulation of steroidogenesis by the bovine corpus luteum. Furthermore, these results suggest that the 5-HT-induced progesterone production is a receptor-mediated event.

Introduction

At the level of the ovary the adrenergic agents adrenaline, noradrenaline and isoproterenol have been shown to stimulate cAMP and progesterone production by luteal tissue from the cow (Condon & Black, 1976; Godkin, Black & Duby, 1977; Milvae, Allia & Hansel, 1983), rat (Harwood, Richert, Dufau & Catt, 1980; Ratner, Sanborn & Weiss, 1980; Norjavaara, Selstam & Ahren, 1982), rabbit (Birnbaumer, Yang, Hunzicker-Dunn, Brockaert & Duran, 1976), and ewe (Jordan, Caffrey & Niswender, 1978). In contrast to the stimulatory effects of catecholamines in the aforementioned species, corpora lutea of the mare (Condon, Ganjam & Kenney, 1979) and woman (Richardson & Masson, 1980; Casper & Cotterell, 1984) appear to be refractory to stimulation by exogenous adrenergic agonists.

Catecholamines act directly on luteal cells, since fibroblasts and erythrocytes separated from ewe luteal cell suspensions did not produce significant amounts of cAMP in response to adrenaline (Jordan et al., 1978). Adrenergic agents are thought to induce cAMP and subsequent progesterone production by binding to specific receptors in the luteal cell membrane (Lefkowitz, 1976). Initial support for this hypothesis is suggested by the finding that propranolol can block the catecholamine-induced increase in luteal cAMP and progesterone production in the cow (Condon & Black, 1976; Godkin et al., 1977), rat (Harwood et al., 1980; Norjavaara et al., 1982) and ewe (Jordan et al., 1978). These results not only suggest specific adrenergic receptors, but also that they are of the β -adrenergic type. In the rat the β -adrenergic stimulation of ovarian cAMP and progesterone has been shown to be mediated by β_2 -

adrenergic receptors (Coleman, Paterson & Somerville, 1979; Ratner, Weiss & Sanborn, 1980; Adashi & Hsueh, 1981; Norjavaara, Roseberg, Gafvels & Selstam, 1984). In the cow and ewe the characterization of a specific β_2 -adrenergic mediation of catecholamine response by luteal cells has not been established. The purpose of this study with cow luteal cells was to determine (a) whether the catecholamine-induced increase in progesterone production is mediated by a β_2 -adrenergic receptor and (b) the role of other biogenic amines on basal and LH-stimulated progesterone production.

Materials and Methods

Materials and drug preparation

Serotonin (5-hydroxytryptamine oxalate salt; 5-HT), di-isoproterenol HCl, acetylcholine chloride, histamine dihydrochloride, salbutamol, γ -amino- α -butyric acid (GABA), melatonin (N-acetyl-5-methoxytryptamine) and di-propranolol HCl were purchased from Sigma Chemical Co., St. Louis, MO. Mianserin HCl and 2-amino-6,7-dihydroxy-1,2,3,4-tetrahydronaphthalene hydrobromide were from Research Biochemicals, Inc., Wayland, MA, and butoxamine HCl was from Burrough's Wellcome Co., Research Triangle Park, NC. Terbutaline was donated by Ciba-Geigy Corp., Summit, NJ, practolol was donated by Imperial Chemical Industries PLC, Macclesfield, Cheshire, U.K. Luteinizing hormone (LH) was supplied by the National Institutes of Health (NIAMDD-bLH-4). Ham's F12 culture medium was purchased from Grand Island Biological Co., Grand Island, NY. The collagenase was Worthington, Type 1, Freehold, NJ, and the N-2-hydroxyethylpiperazine-N-2-ethanesulphonic acid (Hepes) was purchased from Calbiochem-Behring, San Diego, CA.

Salbutamol and terbutaline are specific β_2 -adrenergic receptor agonists, while isoproterenol is a nonspecific β -agonist, activating both subtypes of β -receptors. Butoxamine is a specific β_2 -adrenergic receptor antagonist and practolol a specific β_1 -antagonist; propranolol is a nonspecific β -adrenergic blocking agent. Mianserin is a nonspecific serotonin receptor antagonist which blocks both 5-HT₁ and 5-HT₂ receptor subtypes and 2-amino-6,7-dihydroxy-1,2,3,4-tetrahydronaphthalene hydrobromide is a nonspecific dopamine agonist. All hormone and drug preparations were prepared in cold 0.05M phosphate-

buffered saline (pH 7.4) before use in each experiment and used at a final concentration of 10^{-7} to 10^{-4} M. In all cases N equals the number of animals tested. Duplicate determinations of all treatments were performed for each corpus luteum.

Experimental procedures

Corpora lutea (CL) were obtained per vaginam from regularly cycling, nonlactating dairy cows on days 10-12 of the oestrous cycle (oestrus=Day 0). The luteal tissue was placed into Ham's F12 culture medium containing 24 mM Hepes (pH 7.4) at 4°C for transport back to the laboratory. The luteal tissue was dissociated using collagenase (2000 U/g tissue) as described by Simmons, Caffrey, Phillips, Abel & Niswender (1976) and modified by Pate & Condon (1982). Cell viability was determined by the trypan blue exclusion method (Tennant, 1964). Incubations were conducted utilizing 2.5×10^5 viable cells/culture tube in a final volume of 1 ml of Ham's F12 culture medium. Luteal cell suspensions were incubated in a Dubnoff metabolic shaker water bath at 37°C starting with a 15 min preincubation period, after which the cells were centrifuged (800 g, 10 min at 4°C) and the medium discarded. The cells were then resuspended in fresh medium and the appropriate treatment solutions added. After the incubation period of 2-h, cell suspensions were centrifuged and the culture medium was collected and stored (-20°C) for subsequent determination of progesterone concentration.

Progesterone present in the culture medium was quantitated by radioimmunoassay of unextracted samples using antiprogestosterone-11-bovine serum albumin (No. 337, Niswender). This antiserum does not cross-react

significantly with any other steroid present in the incubation medium (Gibori, Antczak, & Rothchild, 1977). The progesterone tracer used was [1,2-³H] progesterone and was purchased from New England Nuclear (Boston, MA). The sensitivity of the assay as determined by the lower 95% confidence limit of the maximum binding in the absence of any unlabeled progesterone was 0.1 ng. The intraassay and interassay coefficients of variability were 5.3% and 13.3%, respectively. All standards were assayed in quadruplicate, and all unknowns were assayed in duplicate.

Statistical analysis

Analysis of differences between treatment means was conducted using one-way analysis of variance followed by the Student-Newman-Keuls mean separation procedure.

Results

The effects of some of the biogenic amines on basal progesterone production are presented in Table 1. Progesterone production was not affected significantly by acetylcholine, histamine, GABA, dopamine or melatonin over the treatment concentrations tested. Similarly, none of these compounds had any effect on LH-stimulated progesterone production ($P>0.05$, Table 2). Terbutaline and salbutamol did not affect the basal or LH-stimulated production of progesterone. LH (100 ng/ml) alone resulted in a large increase in progesterone production ($P<0.01$).

The addition of serotonin at concentrations from 10^{-6} to 10^{-4} M stimulated the production of progesterone by luteal cells ($P<0.05$, Text-fig. 1). Additionally, the 5-HT induced progesterone response could be blocked by the addition of mianserin (10^{-5} M, $P<0.05$). The 5-HT stimulation of progesterone production was significantly less than that observed for LH alone and there was no additive effect of 5-HT combined with LH. Isoproterenol, at concentrations from 10^{-7} to 10^{-4} M, also resulted in stimulation of progesterone production ($P<0.05$, Text-fig. 2). The combination of isoproterenol plus LH or isoproterenol plus 5-HT (10^{-5} M) did not result in a significant elevation of progesterone above that observed with LH or isoproterenol, respectively ($P>0.05$). The effect of various β -adrenergic receptor antagonists and mianserin upon isoproterenol-stimulated progesterone is presented in Text-fig. 3. The isoproterenol-induced response could not be inhibited by butoxamine (10^{-5} M) or practolol (10^{-5} M) at any of the isoproterenol concentrations tested ($P>0.05$). Similarly, mianserin (10^{-5} M) did not block progesterone production by isoproterenol ($P>0.05$). In contrast,

propranolol (10^{-5} M) did inhibit the isoproterenol-induced response at an isoproterenol concentration of 10^{-7} to 10^{-5} M ($P < 0.05$). None of the antagonists used in these studies significantly affected either basal or LH-stimulated progesterone production (data not shown).

Discussion

These results demonstrate a possible role for serotonin in the stimulation of steroidogenesis by the bovine CL. That the serotonin-induced stimulation of progesterone production could be inhibited by a serotonin receptor antagonist suggests its action to be a receptor-mediated event. A preliminary experiment with C. E. Rexroad, Jr. (USDA, Beltsville) has shown serotonin to be present within the bovine CL at a relatively high concentration (152.7 ± 40.19 ng/g). In the pineal gland of the cow, serotonin concentrations range from 1000 to 2000 ng/g depending upon the time of year (Philo & Reiter, 1980). In the rat, Clausell & Soliman (1978) have reported 5-HT to be present in ovarian tissue: the concentration of 5-HT was approximately 900 ng/g during both dioestrus and metoestrus with a significant increase to 1300 ng/g occurring at oestrus and a role for ovarian 5-HT in the process of ovulation was proposed.

Possible sources of luteal 5-HT include mast cells, blood platelets and de novo synthesis. In this laboratory, histological determination of mast cells using Giemsa, toluidene blue and methylene blue staining techniques has not conclusively demonstrated their presence in luteal tissue of cows. It is possible that serotonin may be working indirectly to stimulate progesterone production by acting on a secondary cell type such as endothelial cells to produce a luteotrophin. Studies are presently underway to evaluate this possibility and to further define the role of serotonin and its metabolites in the stimulation of steroidogenesis by the bovine CL.

In vitro studies using human CL slices showed that melatonin

stimulated progesterone production in a dose-dependent manner while serotonin and N-acetyl serotonin had no effect on progesterone production (MacPhee, Cole, & Rice, 1975). In cultured rat granulosa cells, melatonin has been reported to increase basal and LH-stimulated progesterone production (Fiske, Parker, Ulmer, Ow & Aziz, 1984). [^3H]-melatonin has been reported to be selectively taken up by the ovaries in the rat and cat in vivo (Wurtman, Axelrod & Potter, 1964). In male rats melatonin and serotonin have been demonstrated to inhibit the in vitro biosynthesis of androgens (Peat & Kinson, 1971, Ellis, 1972). In our studies utilizing dissociated bovine luteal cells melatonin was unable to alter either basal or LH-stimulated progesterone production.

A possible role for GABA in the regulation of ovarian function has been suggested due to the relatively high concentration of this compound in rat ovarian tissue (Del Rio & Caballero, 1980). Shaeffer & Hsueh (1982) have reported GABA and glutamate decarboxylase (GAD), the enzyme responsible for GABA synthesis, to be present in whole ovarian homogenates but not in enriched ovarian granulosa cells. Additionally, these authors have demonstrated specific binding sites for GABA in the rat ovary, with the majority of these sites being associated with granulosa cells. The testing of GABA in our short-term incubation system does not support a role for GABA in the control of steroidogenesis by the bovine CL.

To our knowledge, there is only a single report in the literature concerning the effects of acetylcholine on luteal function. In that study, acetylcholine and carbachol (a synthetic cholinergic agent) at concentrations up to 10^{-4}M did not effect basal or LH-stimulated progesterone production by human luteal tissue in either a short-term (4

h) or long-term (10 day) culture system (Casper & Cotterell, 1984). There are no data concerning the effects of histamine and dopamine on luteal steroidogenesis. However, histamine has been reported to induce the production of cortisol and corticosterone by dissociated adrenal cells of the dog (Hirose, Matsumoto & Aikawa, 1978). In our studies we have been unable to demonstrate any significant influence of acetylcholine, histamine or dopamine in controlling luteal steroidogenesis.

The Isoproterenol stimulation of progesterone reported in this study is consistent with previous results reported for bovine luteal slices (Condon & Black; 1976, Godkin et al., 1977) and dissociated bovine luteal cells (Milvae et al., 1983). Neither terbutaline nor salbutamol, the two β_2 -adrenergic agonists used in the present study, significantly affected progesterone production by bovine luteal cells. These results differ from those reported for rat luteal tissue (Ratner et al., 1980; Norjavaara et al., 1984) and FSH-primed rat granulosa cells (Adashi & Hsueh, 1981). Additionally, Isoproterenol was reported to stimulate cAMP by rat luteal tissue, and this response could not be blocked by a β_1 - or a β_2 -adrenergic receptor antagonist, although it could be blocked by propranolol. Similar results have been found in this study for progesterone production by bovine luteal cells. The Isoproterenol stimulation of progesterone production could not be blocked by butoxamine or practolol, but was inhibited by propranolol. The effects of Isoproterenol either alone or in combination with these antagonists, on progesterone production by rat luteal tissue was not reported by Ratner et al. (1980). It may be that the β -adrenergic agonists used in these studies lack the specificity and affinity for β -adrenergic

receptors on bovine luteal cell membranes. In the rat, salbutamol and terbutaline have been shown to be partial agonists and their affinity varies between tissue types (Coleman et al., 1979; Adashi & Hsueh, 1981; Laszlovszky & Erdo, 1983; Norjavaara et al., 1984). As suggested by Coleman, Paterson & Somerville (1979) it is also possible that these β -adrenergic agonists and antagonists are metabolized more readily than isoproterenol or propranolol thus reducing their biological activity. Since neither, β_1 -, nor β_2 -adrenergic antagonists blocked the isoproterenol stimulation of progesterone but propranolol, an adrenergic antagonist which blocks both receptor subtypes, was able to, these results may suggest a mixed population of β -adrenergic receptors on bovine luteal cells.

Scientific Contribution Number 1348 from the New Hampshire Agricultural Experiment station. We thank Dr. G.D. Niswender for the gift of progesterone antiserum, and Dr. Salvatore Raitl of the National Hormone and Pituitary Program of the NIADDK-NIH for the LH used in this study. This research was a contribution to Northeast Regional Project NE-72.

References

- Adashi, E.Y. & Hsueh, A.J.W. (1981) Stimulation of β_2 -adrenergic responsiveness by follicle-stimulating hormone in rat granulosa cells *in vitro* and *in vivo*. *Endocrinology* 108, 2170-2178.
- Birnbaumer, L., Yang, P. C., Hunzicker-Dunn, M., Brockaert, J. & Duran, J. M. (1976) Adenyl cyclase activities in ovarian tissues. I. Homogenization and conditions of assay in Graafian follicles and corpora lutea of rabbits, rats and pigs: regulation by ATP, and some comparative properties. *Endocrinology* 99, 163-184.
- Casper, R. F. & Cotterell, M. A. (1984) The effects of adrenergic and cholinergic agents on progesterone production by human corpus luteum *in vitro*. *Am. J. Obstet. Gynecol.* 148, 663-669.
- Clausell, D. E. & Soliman, K F. A. (1978) Ovarian serotonin content in relation to ovulation. *Experientia* 34, 410-411.
- Coleman, A.J., Paterson, D.S. & Somerville, A.R. (1979a) The adrenergic receptor of rat corpus luteum membranes. *Biochem. Pharmacol.* 28, 1003-1010.
- Coleman, A.J., Paterson, D.S. & Somerville, A.R. (1979b) Factors controlling the selectivity of β blocking drugs. *Biochem. Pharmacol.* 28, 1011-1013.
- Condon, W. A. & Black D. L. (1976) Catecholamine-induced stimulation of progesterone by the bovine corpus luteum *in vitro*. *Biol. Reprod.* 15, 573-578.
- Condon, W. A. Ganjam, V. K. & Kennerly, R. M. (1979) Catecholamines and equine luteal progestagens. *J. Reprod. Fert., Suppl.* 27, 199-203.
- Del Rio, R. M. & Caballero, A. L. (1980) Presence of γ -aminobutyric acid in rat ovary. *J. Neurochem.* 34, 1584-1586.
- Ellis, L. C. (1972) Inhibition of rat testicular androgen synthesis *in vitro* by melatonin and serotonin. *Endocrinology* 90, 17-28.
- Fiske, V. M., Parker, K. L. Ulmer, R. A., Ow, C. H. & Aziz, N. (1984) Effect of melatonin alone or in combination with human chorionic gonadotropin or ovine luteinizing hormone on the *in vitro* secretion of estrogens or progesterone by granulosa cells of rats. *Endocrinology* 114, 407-410.
- Gibori, G., Antczak, E. & Rothchild, I. (1977) The role of estrogen in the regulation of luteal progesterone secretion in the rat after day 12 of pregnancy. *Endocrinology* 100, 1483-1495.
- Godkin, J. D., Black, D. L. & Duby, R. T. (1977) Stimulation of cyclic AMP and progesterone synthesis by LH, PGE_2 and isoproterenol in the bovine CL *in vitro*. *Biol. Reprod.* 17, 514-518.
- Harwood, J. P., Richert, N. D., Dufau, M. L. & Catt, K. J. (1980)

Gonadotropin-induced desensitization of epinephrine action in the luteinized rat ovary. Endocrinology 107, 280-288.

Hirose, T., Matsumoto, I. & Aikawa, T. (1978) Direct effect of histamine on cortisol and corticosterone production by isolated dog adrenal cells. J. Endocr. 76, 371-372.

Jordan, A. W. III, Caffrey, J. L. & Niswender, G. D. (1978) Catecholamine-induced stimulation of progesterone and adenosine 3', 5'-monophosphate production by dispersed ovine luteal cells. Endocrinology 103, 385-392.

Laszlovszky, I. & Erdo, S.L. (1983) Characterization of β -adrenergic receptors in rat ovary. Eur. J. Pharmacol. 96, 101-104.

Lefkowitz, R. J. (1976) β -adrenergic receptors: recognition and regulation. N. Engl. J. Med. 295, 323-328.

MacPhee, A. A., Cole, F. E. & Rice, B. F. (1975) The effect of melatonin on steroidogenesis by the human ovary *in vitro*. J. clin. Endocr. Metab. 40, 688-696.

Milvae, R. A., Ailla, H.W. & Hansel, W. (1983) Methylation in bovine luteal cells as a regulatory of luteinizing hormone action. Biol. Reprod. 29, 849-855.

Norjavaara, E., Rosberg, S., Gafvels, M. & Selstam, G. (1984) β -adrenergic receptor concentration in corpora lutea of different ages obtained from pregnant mare serum gonadotropin-treated rats. Endocrinology 114, 2154-2159.

Norjavaara, E., Selstam, G. & Ahren, K. (1982) Catecholamine stimulation of cyclic AMP and progesterone production in rat corpora lutea of different ages. Acta endocr., Copenh. 100, 613-622.

Pate, J. L. & Condon, W. A. (1982) Effects of serum and lipoproteins on steroidogenesis in cultured bovine luteal cells. Molec. cell. Endocr. 28, 551-562.

Peat, F. & Kinson, G. A. (1971) Testicular steroidogenesis *in vitro* in the rat in response to blinding, pinealectomy and to the addition of melatonin. Steroids 17, 251-264.

Philo, R. & Reiter, R. J. (1980) A circannual rhythm in bovine pineal serotonin. Experientia 36, 664-665.

Ratner, A., Sanborn, C. R. & Weiss, G. K. (1980) β -adrenergic stimulation of cAMP and progesterone in rat ovarian tissue. Am. J. Physiol. 239, E139-E143.

Ratner, A., Weiss, G. K. & Sanborn, C. R. (1980) Stimulation by β_2 -adrenergic receptors of the production of cyclic AMP and progesterone in rat ovarian tissue. J. Endocr. 87, 123-129.

Richardson, M. C. & Masson, G. M. (1980) Progesterone production by

- dispersed cells from human corpus luteum: stimulation by gonadotrophins prostaglandin $F_{2\alpha}$; lack of response to adrenaline and isoprenaline. J. Endocr. 87, 247-254.
- Shaeffer, J. M. & Hsueh, A. J. W. (1982) Identification of gamma-aminobutyric acid and its binding sites in the rat ovary. Life Sci. 30, 1599-1604.
- Simmons, K. R., Caffrey, J. L., Phillips, J. L., Abel, J. H. Jr. & Niswender, G. D. (1976) A simple method for preparing suspensions of luteal cells. Proc. Soc. exp. Biol. Med. 152, 366-371.
- Tennant, J. R. (1964) Evaluation of the trypan blue technique for determination of cell viability. Transplantation 2, 685-694.
- Wurtman, R. J., Axelrod, J. & Potter, L. T. (1964) The uptake of H^3 -melatonin in endocrine and nervous tissues and the effects of constant light exposure. J. Pharmacol. exp. Ther. 143, 314-321.

Table 1. Effect of biogenic amines on progesterone production (ng/ml culture medium) by dissociated luteal cells in a 2-h incubation.

Treatment	N	0	Treatment concentration			
			10 ⁻⁴ M	10 ⁻⁵ M	10 ⁻⁶ M	10 ⁻⁷ M
Control	14	42.6 ± 3.22				
LH (100 ng/ml)	14	188.3 ± 14.94*				
Acetylcholine	(N)		57.3 ± 6.26 ⁽⁵⁾	51.5 ± 5.04 ⁽⁸⁾	59.1 ± 7.28 ⁽⁸⁾	57.1 ± 4.73 ⁽⁷⁾
Histamine	4		61.7 ± 13.67	46.9 ± 8.98	44.1 ± 6.80	44.2 ± 5.90
γ-Aminobutyric Acid	4			45.5 ± 4.90	53.0 ± 13.03	47.8 ± 10.86
Dopamine Agonist	4			47.6 ± 3.65	43.4 ± 6.62	44.8 ± 8.07
Melatonin	4			54.9 ± 7.92	54.5 ± 4.76	43.6 ± 4.19
Terbutaline	3		65.3 ± 8.20	58.8 ± 16.49	53.6 ± 7.61	41.5 ± 9.38
Salbutamol	6		69.4 ± 5.66	61.7 ± 5.53	63.8 ± 4.41	49.7 ± 3.28

Values are the mean ± s.e.m.

*P<0.01 compared with control value

Table 2. Effect of biogenic amines on LH-stimulated progesterone production (ng/ml culture medium) by dissociated luteal cells in a 2-h incubation.

Treatment	N	Treatment Concentration				
		0	10^{-4}M	10^{-5}M	10^{-6}M	10^{-7}M
Control	14	42.6 ± 3.22				
LH (100 ng)	14	$188.3 \pm 14.94^*$				
Acetylcholine	(5)		191.1 ± 36.96	211.1 ± 27.64	221.1 ± 29.99	197.1 ± 26.08
Histamine	3		156.0 ± 24.29	149.2 ± 21.24	153.4 ± 22.59	149.6 ± 25.19
γ -Aminobutyric Acid	3			168.4 ± 11.85	176.6 ± 13.96	214.8 ± 20.03
Dopamine Agonist	3			198.6 ± 45.88	188.8 ± 46.01	179.9 ± 42.02
Melatonin	3			212.7 ± 10.25	216.1 ± 9.00	212.2 ± 17.16
Terbutaline	3		186.6 ± 18.76	162.8 ± 27.04	148.1 ± 31.05	140.3 ± 24.11
Salbutamol	3		200.6 ± 51.96	203.8 ± 36.28	202.5 ± 48.06	197.7 ± 26.06

Values are the mean \pm s.e.m.


* $P < 0.01$ compared with control value


Figure Legends


Text-fig. 1. Effect of serotonin on progesterone production by dissociated luteal cells. Data are expressed as the mean \pm s.e.m. $N = 6(10^{-4}M)$ and $8(10^{-7}, 10^{-6}$ and $10^{-5}M)$ for 5-HT, $N = 3$ for 5-HT + LH. $N = 6$ for 5-HT + Mianserin (mian) and $N = 14$ for control (C) and LH. Bars with different superscript letters are significantly different ($P < 0.01$, LH and 5-HT + LH; $P < 0.05$, 5-HT and 5-HT + mianserin)

Text-fig. 2. Effect of Isoproterenol on luteal cell progesterone production. Data are expressed as the mean \pm s.e.m. $N = 5(10^{-4}M)$ and $6(10^{-7}, 10^{-6}$ and $10^{-5}M)$ for Isoproterenol (Iso). $N = 3$ for Isoproterenol + LH and Isoproterenol + 5-HT, $N = 14$ for control (C) and LH. Bars with different superscript letters are significantly different ($P < 0.01$, LH and ISO + LH; $P < 0.05$ for Isoproterenol and Isoproterenol + 5-HT).

Text-fig. 3. Effect of β -adrenergic antagonists and mianserin on isoproterenol-stimulated progesterone production. Data are expressed as the mean \pm s.e.m. $N = 4$ for Isoproterenol (iso) + butoxamine (butox) and isoproterenol + mianserin (mian), $N = 3$ for isoproterenol + practolol (pract) and isoproterenol + propranolol (propran). Bars with different superscript letters are significantly different from isoproterenol alone (—) ($P < 0.05$, for isoproterenol + propranolol).







EXPERIMENT TWO

Mechanisms involved in the action of serotonin-induced stimulation
of progesterone production by bovine luteal cells in vitro

P. J. Battista, C. E. Rexroad, Jr.¹

and W. A. Condon

Department of Animal and Nutritional Sciences

The University of New Hampshire

Durham, New Hampshire 03824 (U.S.A.)

and

U.S. Department of Agriculture¹

Beltsville, Maryland 20705 (U.S.A.)

Accepted for publication in the Journal of
Molecular and Cellular Endocrinology

Summary

Serotonin (5-HT) induced stimulation of progesterone (P_4) production by bovine luteal cells was characterized with respect to a) the receptor subtype mediating this response, b) the steroidogenic response to 5-HT metabolites, c) the role of adenylate cyclase and d) the 5-HT concentration of bovine luteal tissue. Addition of 5-HT ($10^{-5}M$) stimulated the production of P_4 ($P<0.05$) and this stimulation was inhibited by the 5-HT antagonist mianserin at a concentration of $10^{-5}M$ ($P<0.05$), but not at a mianserin concentration of $10^{-7}M$. Additionally, the response to 5-HT could not be inhibited by ketanserin ($10^{-5}M$), a 5-HT₂ receptor antagonist. Incubation of luteal cells with a specific 5-HT₁ agonist, DPAT ($10^{-4}M$), stimulated the production of P_4 ($P<0.05$) and this response could not be blocked by mianserin at $10^{-7}M$ or by ketanserin, but was inhibited by mianserin at $10^{-5}M$ ($P<0.05$). The addition of the 5-HT metabolite 5-methoxytryptamine (5-MTA) stimulated P_4 production ($P<0.05$) and this response could be inhibited by mianserin ($10^{-5}M$, $P<0.05$). Neither N-acetyl-5-HT nor 5-methoxytryptophan significantly affected P_4 production. The addition of the phosphodiesterase inhibitor 3-isobutyl-methylxanthine (IBMX, 0.1mM), potentiated the effects of 5-HT and DPAT ($P<0.05$), but this effect was additive rather than synergistic. In contrast, the addition of luteinizing hormone (10 ng/ml) plus IBMX resulted in a significant synergistic response ($P<0.05$). Analysis of luteal tissue by high performance liquid chromatography showed the concentration of 5-HT to vary with the stage of corpora lutea development, with the highest

concentration being present in early (days 4-6) luteal tissue ($P < 0.05$). These results demonstrate the presence of 5-HT within bovine luteal tissue and suggest that the 5-HT and 5-MTA stimulation of P_4 are mediated by a 5-HT₁ receptor which may not be associated with adenylate cyclase.

Introduction

Previous work from this laboratory has shown that serotonin (5-HT) stimulates the production of progesterone (P_4) by dissociated bovine luteal cells in short-term incubation (Battista and Condon, 1986). Additionally, this response could be inhibited by the addition of a 5-HT receptor antagonist, suggesting its action is receptor-mediated. Radioligand binding studies have revealed at least two distinct classes of recognition sites for 5-HT within the central nervous system (Peroutka and Snyder, 1979a). The 5-HT₁ receptor subtype displays a nM affinity for 5-HT and 5-HT agonists and a μ M affinity for antagonists. The 5-HT₂ receptor site has the opposite pharmacological profile with a nM affinity for antagonists and a μ M affinity for 5-HT agonists (Bennett and Snyder, 1976; Leysen et al., 1978; Peroutka and Snyder, 1979a). The mechanism of signal transduction for the 5-HT₁ receptor has been suggested to involve the adenylate cyclase system (Peroutka et al., 1979b, 1981; Barbaccia et al., 1983) while the 5-HT₂ receptor has been found to be associated with phosphatidylinositol hydrolysis and possible protein kinase C activation (Berridge et al., 1982; Brown et al., 1984; Conn and Sanders-Bush, 1984; Leysen et al., 1984).

The present study further characterized the 5-HT-induced stimulation of P_4 with respect to a) the 5-HT receptor subtype mediating this response, b) the steroidogenic response to 5-HT metabolites and c) the ability of phosphodiesterase inhibition to potentiate the 5-HT response. Additionally, 5-HT concentration in bovine luteal tissue was determined at different stages of luteal development.

Materials and Methods

Materials, drug description and preparation

Serotonin (5-hydroxytryptamine oxalate salt; 5-HT), N-acetyl-5-hydroxytryptamine (N-acetyl-5-HT), 5-methoxy-dl-tryptophan (5-MTP), 5-methoxytryptamine HCl (5-MTA) and 3-isobutyl-methylxanthine (IBMX) were purchased from Sigma Chemical Co. (St. Louis, MO.). Mianserin HCl and (\pm)-8-hydroxydiethylaminotetralin HBr (DPAT) were from Research Biochemicals, Inc. (Wayland, MA.) and ketanserin was donated by Janssen Pharmaceutica, Inc. (Beerse, Belgium). Luteinizing hormone (LH) was supplied by the National Institutes of Health (NIAMMD-bLH-4). Ham's F12 culture medium was purchased from Grand Island Biological Co. (Grand Island, NY.). The collagenase was Worthington, Type 1, (Freehold, NJ) and N-2-hydroxyethylpiperazine-N-2-ethanesulfonic acid (Hepes) was purchased from Calbiochem-Behring (San Diego, CA.).

Mianserin is a nonspecific serotonin antagonist which blocks both 5-HT₁ and 5-HT₂ receptor subtypes depending on concentration (Bennett and Snyder, 1976; Peroutka and Snyder, 1979). Ketanserin is a specific 5-HT₂ receptor antagonist (Leysen et al., 1981, 1982) and DPAT is a specific 5-HT₁ receptor agonist (Hjorth et al., 1982; Middlemiss and Fozard, 1982). IBMX is a phosphodiesterase inhibitor which acts to elevate intracellular concentrations of cAMP (Butcher and Sutherland, 1962). All hormones and drugs were prepared in cold 0.05M phosphate-buffered saline (pH 7.2) prior to use in each experiment. In all cases n equals the number of animals tested and duplicate determinations of all treatments were performed for each corpus luteum.

Tissue dissociation and incubation

Corpora lutea (CL) were obtained per vaginum from regularly cycling, nonlactating dairy cows on days 10-12 of the estrous cycle (estrus=Day 0). Luteal tissue was placed into Ham's F12 culture medium containing 24mM Hepes (pH 7.4) at 4°C for transport to the laboratory. Luteal tissue was dissociated with collagenase (2000 u/g tissue) and cell number and viability determined as described by Pate and Condon (1982). Incubations were conducted utilizing 2.5×10^5 viable cells in a final volume of 1ml of Ham's F12 culture medium and incubated at 37°C for 2h as previously described (Battista and Condon, 1986).

Progesterone in the incubation medium was quantitated by radioimmunoassay of unextracted samples using antiprogestosterone-11-bovine serum albumin (No. 337, Niswender). The progesterone tracer used was [1,2-³H] progesterone (New England Nuclear, Boston, MA). The sensitivity of the assay, as determined by the lower 95% confidence limit of the maximum binding in the absence of any unlabeled progesterone, was 0.1ng. The intra-assay and inter-assay coefficients of variability were 6.4% and 11.7%, respectively. All standards were assayed in quadruplicate, and all unknowns were assayed in duplicate.

Experimental procedures

To determine the 5-HT receptor subtype involved in initiating the 5-HT-induced stimulation of P_4 , luteal cell suspensions were incubated with 5-HT (10^{-5} M) or DPAT (10^{-4} M) either alone or in combination with mianserin (10^{-7} M and 10^{-5}) or ketanserin (10^{-5} M). Control tubes (no additions) were included in each experiment. To evaluate the steroidogenic potential of serotonin metabolites, N-acetyl-5-HT (10^{-5} - 10^{-4} M), 5-MTP (10^{-5} - 10^{-4} M) and 5-MTA (10^{-7} - 10^{-4} M) were added to

Incubations and their ability to modulate P_4 production evaluated. To determine if an increase in intracellular cAMP concentration could potentiate the response to 5-HT, luteal cells were incubated with the following treatments in the presence and absence of 0.1mM IBMX: 5-HT (10^{-5} M), DPAT (10^{-4} M), 5-HT + mianserin (10^{-7} and 10^{-5} M), 5-HT + ketanserin (10^{-5} M) and IBMX alone (control).

For the determination of 5-HT concentration, CL were removed per vagium at various stages of the estrous cycle: days 4-6 (early) days 10-12 (mid) and days 15-16 (late). Upon removal, tissue was rinsed with Ham's F12 then cut in half and a cross-sectional slice obtained. The tissue slice was rinsed again and immediately placed on dry ice for transport to the laboratory. Tissue was stored in liquid nitrogen (liquid phase) prior to analysis. Tissue concentrations of 5-HT were determined using high performance liquid chromatography (HPLC) as described by Mayer and Shoup (1983). Briefly, luteal tissue was homogenized in 5% trichloroacetic acid. Dihydroxybenzylamine was added to the tissue prior to homogenization to serve as an internal standard. The homogenate was filtered through a 0.22 micron filter prior to injecting 50 μ l aliquots onto an HPLC column. The 5-HT was identified by its retention time compared to an external standard and quantitated against the external standard. The internal standard was used to account for recovery. Recovery from the original homogenate of the internal standard was $.90 \pm .02$ (mean \pm SEM, $n=18$).

Statistical analysis

Analysis of differences between treatment means was conducted using one-way analysis of variance followed by the Student-Newman-Keuls mean separation procedure. Two-way analysis of variance was used to test for

a significant interaction in those experiments involving the addition of IBMX.

Results

Incubation of luteal cell suspensions with 5-HT (10^{-5} M) resulted in a significant stimulation of P_4 production ($P < 0.05$, Fig. 1). The addition of mianserin at 10^{-5} M but not at 10^{-7} M, inhibited the 5-HT-induced production of P_4 ($P < 0.05$). Coincubation of 5-HT with ketanserin (10^{-5} M) did not block the steroidogenic action of 5-HT ($P > 0.05$). Addition of DPAT (10^{-4} M) stimulated the production of P_4 ($P < 0.05$, Fig. 2) and this response could not be inhibited by mianserin (10^{-7} M) or by ketanserin (10^{-5} M) ($P > 0.05$) but was blocked by mianserin at 10^{-5} M ($P < 0.05$). The combination of 5-HT + DPAT did not result in a further increase in P_4 production above that observed with 5-HT alone (Figs. 1 and 2) but the response to 5-HT + DPAT was greater than DPAT alone ($P < 0.05$, Fig. 2).

The effects of 5-HT metabolites on P_4 production are presented in Table 1. Addition of 5-MTA at concentrations from 10^{-6} to 10^{-4} M resulted in a dose-dependent stimulation of P_4 production ($P < 0.05$). Mianserin at 10^{-5} M inhibited the 5-MTA response ($P < 0.05$). Neither N-acetyl-5-HT nor 5-MTP significantly affected P_4 production at either of the concentrations tested.

Addition of 0.1mM IBMX to the incubation medium increased basal P_4 production (47.4 ± 1.96 vs 103.5 ± 5.42 for control and IBMX, respectively, $P < 0.05$, Fig. 3). This treatment also significantly potentiated the effects of 5-HT and DPAT, and the potentiation of 5-HT could be blocked by the addition of mianserin at 10^{-5} M ($P < 0.05$) but not at a mianserin concentration of 10^{-7} M or by ketanserin ($P > 0.05$, Fig. 3). The response to 5-HT was greater than that found for DPAT ($P < 0.05$). The

potentiation observed when 5-HT or DPAT was colncubated with IBMX was additive rather than synergistic. In contrast, incubation of luteal cells with LH (10ng/ml) plus IBMX resulted in a significant synergistic response (255.5 ± 15.69 vs 358.7 ± 18.48 for LH and LH + IBMX, respectively, $n=6$, $P<0.05$). Neither mianserin nor ketanserin significantly affected basal P_4 production (51.1 ± 4.76 and 47.8 ± 3.28 for mianserin and ketanserin, respectively, $n=6$, $P>0.05$).

The results of the HPLC analysis of luteal 5-HT concentration are presented in Fig. 4. Concentration of 5-HT was found to vary significantly with the stage of the estrous cycle ($P<0.05$). 5-HT was in highest concentration in luteal tissue obtained during the early luteal phase and was found to decline as CL development proceeded.

Discussion

The ability of 5-HT to stimulate P_4 production by bovine luteal cells is consistent with our previously reported results (Battista and Condon, 1986). The inability of low molar concentrations of mianserin and the specific 5-HT₂ receptor antagonist ketanserin, to inhibit the action of 5-HT suggests the 5-HT response is mediated by the 5-HT₁ receptor subtype. This conclusion is strengthened by the finding that DPAT, a specific 5-HT₁ agonist, could also significantly stimulate the production of progesterone. Additionally, the action of DPAT could not be inhibited by low molar concentrations of mianserin or by ketanserin but was inhibited by higher concentrations of mianserin.

The addition of the 5-HT metabolite 5-MTA resulted in a dose-dependent stimulation of P_4 similar to that previously reported for 5-HT (Battista and Condon, 1986). The increase in P_4 production in response to 10^{-5} M of 5-HT and 5-MTA was similar. These results suggest 5-MTA is as active as 5-HT in vitro. 5-MTA appears to be acting through the same receptor as 5-HT, since the action of 5-MTA could also be blocked by mianserin. Additional support for this conclusion is drawn from the observations of Bennett and Snyder (1976) who reported 5-MTA was effective in the displacement of [³H] 5-HT binding from cerebral cortex membranes. These authors also reported 5-MTA to be more potent in the displacement of binding from the 5-HT₁ receptor subtype. These results lend further support for a 5-HT₁ receptor site on bovine luteal cell membranes.

In vitro studies utilizing human CL slices showed melatonin stimulated P_4 production, while 5-HT and N-acetyl-5-HT were ineffective

In regulating the production of P_4 (MacPhee et al., 1975). 5-HT has been demonstrated to stimulate hydrocortisone secretion in perfused adrenal gland preparations of the dog (Verdesca et al., 1961). In male rats melatonin and 5-HT inhibit the in vitro biosynthesis of androgens (Peat and Kinson, 1971; Ellis, 1972). We have previously reported that melatonin was unable to alter basal or LH-stimulated P_4 by dissociated bovine luteal cells (Battista and Condon, 1986). In the present study, neither N-acetyl-5-HT nor 5-MTP were effective in regulating steroidogenesis. The inability of N-acetyl-5-HT or melatonin to stimulate P_4 production suggests that the action of 5-HT is not due to its conversions to these metabolites.

The ability of the phosphodiesterase inhibitor IBMX to increase basal and LH-stimulated P_4 production demonstrates our short-term culture system is responsive to increased intracellular concentrations of cAMP. The inability of IBMX to potentiate the response to 5-HT in a synergistic manner may suggest a mechanism of signal transduction not associated with the adenylate cyclase system. These results differ from the 5-HT₁ receptor found within the central nervous system, which has been suggested to involve adenylate cyclase (Peroutka et al., 1979b, 1981; Barbaccia et al., 1983). It is possible that the 5-HT₁ receptor associated with adenylate cyclase suggested by Barbaccia et al. (1983) and the 5-HT₁ receptor found on bovine luteal cell membranes represent different subclasses of the 5-HT₁ receptor which elicit responses through different mechanisms. It is also likely that the 5-HT₁ receptor found on bovine luteal cells is of the 5-HT_{1A} subclass since DPAT has been reported to have a greater affinity for the 5-HT_{1A} receptor site (Middlemiss and Fozard, 1983; Hoyer et al., 1985).

The results of the HPLC analysis demonstrate the presence of 5-HT within bovine luteal tissue and show that the concentration of 5-HT varies with respect to the stage of CL development. In the rat, Clausell and Soliman (1978) have reported 5-HT to be present in whole ovarian homogenates. The concentration of 5-HT was found to vary with the stage of the cycle, with the highest concentrations occurring at estrus. Additionally, 5-HT has been reported in rat testicular tissue (Zieher et al., 1971; Ellis, 1972). Possible sources of luteal 5-HT include mast cells, blood platelets, storage of peripherally circulating 5-HT or de novo synthesis. Histological determination of mast cells using Giemsa, toluidene blue and methylene blue staining techniques has failed to demonstrate the presence of mast cells in either early or mid-cycle luteal tissue (unpublished observations). 5-HT present in luteal tissue may be derived from peripherally circulating 5-HT similar to that reported for rat testicular (Ellis et al., 1972) and adrenal (Verhofstad and Jonsson, 1983) tissues. Although the exact mechanism whereby 5-HT regulates P_4 production is unknown, the ability of 5-HT to stimulate luteal steroidogenesis and the presence of 5-HT within luteal tissue may suggest a physiological role for 5-HT as an intraovarian stimulator of luteal steroidogenesis.

The authors gratefully acknowledge M. Burg and A. Powell for technical assistance and D. Stranger for manuscript preparation. We also thank Dr. G.D. Niswender for the progesterone antiserum and Dr. S. Raiti for the LH used in these studies.

References

- Barbaccia ML, Brunello N, Chuang DM, Costa E, 1983. Serotonin-elicited amplification of adenylate cyclase activity in hippocampal membranes from adult rat. *J Neurochem* 40: 1671-79
- Battista PJ, Condon WA, 1986. Serotonin-induced stimulation of progesterone production by cow luteal cells in vitro. *J Reprod Fert* 76: 231-38
- Bennett JP, Snyder SH, 1976. Serotonin and lysergic acid diethylamide binding in rat brain membranes: relationship to postsynaptic serotonin receptors. *Molec Pharmacol* 12: 373-89
- Berridge MJ, Downes CP, Hanley MR, 1982. Lithium amplifies agonist-dependent phosphatidylinositol responses in brain and salivary gland. *Biochem J* 206: 587-95
- Blinks JR, Olson CB, Jewell BR, Braveny P, 1972. Influence of caffeine and other methylxanthines on mechanical properties of isolated mammalian heart muscle. *Circ Res* 30: 367-92
- Brown E, Kendall DA, Nahorski SR, 1984. Inositol phospholipid hydrolysis in rat cerebral cortical slices: I. receptor characterization. *J Neurochem* 42: 1379-87
- Butcher RW, Sutherland EW, 1962. Adenosine 3', 5'-phosphate in biological materials. I. Purification and properties of cyclic 3', 5'-nucleotide phosphodiesterase and use of this enzyme to characterize adenosine 3', 5'-phosphate in human urine. *J Biol Chem* 237:1244-50
- Clausell DE, Soliman KFA, 1978. Ovarian serotonin content in relation to ovulation. *Experientia* 34: 410-11
- Conn PJ, Sanders-Bush E, 1984. Selective 5-HT-2 antagonists inhibit serotonin stimulated phosphatidylinositol metabolism in cerebral cortex. *Neuropharmacol* 23: 993-96
- Ellis LC, 1972. Inhibition of rat testicular androgen synthesis in vitro by melatonin and serotonin. *Endocrinology* 90: 17-28
- Ellis LC, Jaussi AW, Baptista MH, Urry RL, 1972. Correlation of age changes in monoamine oxidase activity and androgen synthesis by rat testicular minced and teased-tubular preparation in vitro. *Endocrinology* 90: 1610-18
- Gore SD, Behrman HR, 1984. Alteration of transmembrane sodium and potassium gradients inhibits the action of luteinizing hormone in the luteal cell. *Endocrinology* 114: 2020-31
- Hjorth S, Carlsson A, Lindberg P, Sanchez D, Wikstrom H, Arvidsson LE, Hackzell U, Nilsson JLG, 1982. 8-Hydroxy-2-(di-n-propylamino)

- tetralin, 8-OH-DPAT, a potent and selective simplified ergot congener with central 5-HT-receptor stimulating activity. *J Neural Transm* 55: 169-88
- Leysen JE, Niemegeers CJE, Tollenaere JP, Laduron PM, 1978. Serotonergic component of neuroleptic receptors. *Nature* 272: 168-71
- Leysen JE, Awouters F, Kennis L, Laduron PM, Vandenberg J, Janssen PAJ, 1981. Receptor binding profile of R41468, a novel antagonist at 5-HT₂ receptors. *Life Sci* 28: 1015-22
- Leysen JE, Niemegeers CJE, Van Nueten JM, Laduron PM, 1982. [³H] Ketanserin (R41468), a selective ³H-ligand for serotonin₂ receptor binding sites. Binding properties, brain distribution and functional role. *Molec Pharmacol* 21: 301-14
- Leysen JE, Courcelles DC, Clerck F, Niemegeers CJE, Van Nueten JM, 1984. Serotonin-S₂ receptor binding sites and functional correlates. *Neuropharmacol* 23: 1493-1501
- MacPhee AA, Cole FE, Rice BF, 1975. The effect of melatonin on steroidogenesis by the human ovary in vitro. *J Clin Endocrinol Metab* 40: 688-96
- Mayer GS, Shorep RE, 1983. Simultaneous multiple electrode liquid chromatographic-electrochemical assay for catecholamines, indoleamines and metabolites in brain tissue. *J Chromatogr* 255: 533-44
- Middlemiss DN, Forzard JR, 1983. 8-Hydroxy-2-(di-n-propylamino)-tetralin discriminates between subtypes of the 5-HT₁ recognition site. *Eur J Pharmacol* 90: 151-53
- Pate JL, Condon WA, 1982. Effects of serum and lipoproteins on steroidogenesis in cultured bovine luteal cells. *Molec cell Endocrinol* 28: 551-62
- Peat F, Kinson GA, 1971. Testicular steroidogenesis in vitro in the rat in response to blinding, pinealectomy and to the addition of melatonin. *Steroids* 17: 251-64
- Peroutka SJ, Snyder SH, 1979a. Multiple serotonin receptors: differential binding of [³H] 5-hydroxytryptamine, [³H] lysergic acid diethylamide and [³H] spiroperidol. *Molec Pharmacol* 16: 687-99
- Peroutka SJ, Lebovitz RM, Snyder SH 1979b. Serotonin receptor binding sites affected differentially by guanine nucleotides. *Molec Pharmacol* 16: 700-08
- Peroutka SJ, Lebovitz RM, Snyder SH, 1981. Two distinct central serotonin receptors with different physiological functions. *Science* 212: 827-29
- Shemesh M, Hansel W, Strauss III JF, 1984. Calcium-dependent, cyclic nucleotide-independent steroidogenesis in the bovine placenta. *Proc Natl Acad Sci* 81: 6403-07

- Verdesca AS, Westermann CD, Crampton RS, Black WC, Nedeljkovic RI, Hilton JC, 1961. Direct adrenocortical stimulatory effect of serotonin. *Am J Physiol* 210: 1065-67
- Verhofstad AAJ, Jonsson G, 1983. Immunohistochemical and neurochemical evidence for the presence of serotonin in the adrenal medulla of the rat. *Neuroscience* 10: 1443-53
- Zieher LM, Debeljuk L, Iturriza F, Mancini RE, 1971. Biogenic amine concentration in testes of rats at different ages. *Endocrinology* 88: 351-54.

Table 1. Effect of serotonin metabolites on P_4 production (ng/ml of incubation medium) by dissociated luteal cells in short-term incubation.

Treatment	n	Treatment Concentration				
		0	$10^{-4}M$	$10^{-5}M$	$10^{-6}M$	$10^{-7}M$
Control	6	43.3 ± 3.14^a				
5-MTA	6		146.0 ± 7.87^b	115.5 ± 10.89^c	93.0 ± 6.98^d	60.7 ± 2.37^a
5-MTA + mianserin ($10^{-5}M$)	4		66.2 ± 2.82^a	51.0 ± 5.51^a	50.9 ± 5.83^a	43.6 ± 4.48^a
N-Acetyl-5-HT	6		48.6 ± 4.17^a	52.0 ± 5.57^a	ND	ND
5-MTP	6		41.8 ± 2.56^a	44.8 ± 2.71^a	ND	ND

Values are the mean \pm SEM

Numbers with different superscript are significantly different ($P < 0.05$).

ND denotes not determined


Figure Legends


Fig 1. Effect of 5-HT antagonists on 5-HT-induced P_4 production. Data are expressed as the mean \pm SEM, $n=9$ for control and 5-HT, $n=6$ for all other treatments. MIA=mlanserin and KET=ketanserin. Bars with different superscripts are significantly different ($P<0.05$).


Fig 2. Effect of the 5-HT₁ agonist DPAT on luteal cell P_4 production. Data are expressed as the mean \pm SEM, $n=9$ for control, $n=7$ for all other treatments. MIA=mlanserin and KET=ketanserin. Bars with different superscripts are significantly different ($P<0.05$).


Fig 3. Effect of IBMX on basal P_4 production and P_4 production induced by 5-HT and DPAT either alone or in combination with 5-HT antagonists. Data are expressed as the mean \pm SEM, $n=6$ for all treatments. Solid line denotes control value (no IBMX) of 47.4 ± 1.96 ng/ml. MIA=mlanserin and KET=ketanserin. Bars with different superscripts are significantly different ($P<0.05$).

Fig 4. 5-HT concentration (ng/g of tissue) of bovine luteal tissue as affected by stage of luteotropism. Data are expressed as the mean \pm SEM, $n=5$ for early, $n=8$ for mid and $n=5$ for late stages of the cycle. Bars with different superscripts are significantly different ($P<0.05$).









EXPERIMENT THREE

A role for alternative pathway catecholamines in the
stimulation of steroidogenesis in cow luteal cells

P. J. Battista and W. A. Condon

Department of Animal and Nutritional Sciences,
The University of New Hampshire
Durham, New Hampshire 03824, U.S.A.

Published in the Journal of Reproduction
and Fertility (1986) 78:275-280.

Summary

Incubation of bovine luteal cells with the alternative pathway catecholamines octopamine, synephrine and deoxyadrenaline at concentrations of 10^{-6} to 10^{-3} M enhanced the production of progesterone ($P < 0.05$). Tyramine did not alter basal progesterone production ($P > 0.05$). Addition of noradrenaline and adrenaline at concentrations of 10^{-4} to 10^{-7} M significantly elevated the production of progesterone ($P < 0.05$). The steroidogenic response to noradrenaline and adrenaline was greater than that for octopamine, synephrine and deoxyadrenaline ($P < 0.05$). Response to both primary (10^{-6} M) and alternative (10^{-4} M) pathway catecholamines was inhibited by propranolol (10^{-5} M, $P < 0.05$) but not by phentolamine (10^{-5} M, $P > 0.05$). These results demonstrate that octopamine, synephrine and deoxyadrenaline can affect steroidogenesis by bovine luteal cells, and their action is mediated by β -adrenergic receptors.

Introduction

The primary catecholamines noradrenaline and adrenaline can stimulate the synthesis in vitro of cAMP and progesterone by luteal tissue from a variety of species (cow: Condon & Black, 1976; Godkin, Black & Duby, 1977; Milvae, Allia & Hansel, 1983; rat: Harwood, Dufau & Catt, 1979; Ratner, Sanborn & Weiss, 1980; Norjavaara, Seistam & Ahren, 1982; rabbit: Birnbaumer, Yang, Hunzicker-Dunn, Brockaert & Duran, 1976; ewe: Jordan, Caffrey & Niswender, 1978). Noradrenaline and adrenaline can interact with both α - and β -adrenergic receptors although their stimulatory effects on luteal steroidogenesis are mediated by β -receptors, since their response can be inhibited by β -adrenergic antagonists but not by α -adrenergic antagonists (Condon & Black, 1976; Jordan et al., 1978; Harwood, Richert, Dufau & Catt, 1980; Ratner et al., 1980; Norjavaara et al. 1982). In contrast, nothing is known about the catecholamines derived through the alternative catecholamine pathway (Fig. 1) with respect to ovarian steroidogenesis. Octopamine and synephrine have been identified within the central nervous system and peripheral tissues in a number of vertebrate species (Kakimoto & Armstrong 1962; Molinoff & Axelrod, 1969, 1972; Saavedra, 1974; Harmar & Horn, 1976). In peripheral tissues octopamine is associated with sympathetic nerve endings (Molinoff & Axelrod, 1969, 1972) where it can be taken-up, synthesized and stored (Fischer, Musacchio, Kopin & Axelrod, 1964; Kopin, Fischer, Musacchio & Horst, 1964; Kopin, Fischer, Musacchio, Horst & Weise, 1965; Baldessarini & Vogt, 1971). Octopamine accumulates in sympathetic nerves after monoamine oxidase inhibition and is released following neural stimulation (Kopin et al., 1964, 1965;

Fischer, Horst & Kopin, 1965; Baldessarini, 1971; Baldessarini & Vogt, 1972). Additionally, octopamine has a subcellular distribution similar to that of noradrenaline (Musacchio, Kopin & Snyder, 1964; Snyder, Michaelson & Musacchio, 1964; Snyder, Glowinski & Axelrod, 1965; Baldessarini & Vogt 1971; Molinoff & Axelrod, 1972), and a physiological role for octopamine as a cotransmitter with noradrenaline has been proposed (Molinoff & Axelrod 1969, 1972; Axelrod & Saavedra, 1977).

The physiological significance of the alternative pathway catecholamines with respect to luteal function is unknown. It was the purpose of this study to examine the ability of the alternative pathway catecholamines tyramine, octopamine, synephrine and deoxyadrenaline to stimulate steroidogenesis by bovine luteal cells.

Materials and Methods

Materials, drug description and preparation

Tyramine HCl, DL-octopamine HCl, DL-synephrine, deoxyadrenaline HCl, L-noradrenaline HCl, DL-adrenaline and DL-propranolol HCl were purchased from Sigma Chemical Co., St. Louis, MO. Phentolamine HCl was donated by Ciba-Geigy Corp., Summit, NJ. Ham's F12 culture medium was purchased from Grand Island Biological Co., Grand Island, NY. The collagenase was Worthington, Type 1, Freehold, NJ and N-2-hydroxyethylpiperazine-N-2-ethanesulphonic acid (Hepes) was purchased from Calbiochem-Behring, San Diego, CA.

Propranolol is a nonspecific β -adrenergic receptor antagonist and phentolamine is a nonspecific α -adrenergic antagonist. All hormones and drugs were prepared in cold 0.05 M phosphate-buffered saline (pH 7.2) before use in each experiment. In all cases N equals the number of animals tested. Duplicate determination of all treatments was performed for each corpus luteum.

Experimental procedures

Corpora lutea (CL) were obtained per vaginam from regularly cycling, nonlactating dairy cows on Days 10-12 of the oestrous cycle (oestrus=Day 0). Luteal tissue was placed into Ham's F12 culture medium containing 24 mM-Hepes (pH 7.4) at 4°C for transport to the laboratory. Luteal tissue was dissociated with collagenase (2000 u/g tissue) and cell number and viability were determined as described by Pate & Condon (1982).

Incubations were conducted utilizing 2.5×10^5 viable cells in a

final volume of 1 ml Ham's F12 culture medium and incubated for 2h at 37°C as previously described (Battista & Condon, 1986). Tyramine, octopamine, synephrine and deoxyadrenaline were added to luteal cell suspensions at a final concentration of 10^{-6} to 10^{-3} M. Noradrenaline and adrenaline were tested at 10^{-7} to 10^{-4} M. For determination of the adrenergic receptor mediating the catecholamine response, noradrenaline and adrenaline (10^{-6} M) or octopamine, synephrine and deoxyadrenaline (10^{-4} M) were added to incubations which had been preincubated for 30 min with propranolol (10^{-5} M) or phentolamine (10^{-5} M).

Progesterone in the incubation medium was quantitated by radioimmunoassay of unextracted samples using antiprogesterone-11-bovine serum albumin (No. 337, Niswender). The progesterone tracer used was [1,2- 3 H]progesterone (New England Nuclear, Boston, MA). The sensitivity of the assay, as determined by the lower 95% confidence limit of the maximum binding in the absence of any unlabelled progesterone, was 0.1 ng. The intra-assay and interassay coefficients of variability were 7.3% and 12.4% respectively. All standards were assayed in quadruplicate, and all unknowns were assayed in duplicate.

Statistical analysis

Analysis of difference between treatment means was conducted using one-way analysis of variance followed by the Student-Newman-Keuls mean separation procedure.

Results

Luteal steroidogenesis in the presence of alternative pathway catecholamines is shown in Fig. 2. Octopamine, synephrine and deoxyadrenaline significantly enhanced progesterone production at concentrations of 10^{-6} to 10^{-3} M ($P < 0.05$). Tyramine was ineffective in influencing the production of progesterone ($P > 0.05$). The response to octopamine, synephrine and deoxyadrenaline was similar within each treatment concentration. Addition of noradrenaline and adrenaline at 10^{-7} to 10^{-4} M resulted in a significant elevation in the production of progesterone ($P < 0.05$, Fig. 3.). Comparison of the response of luteal cells to primary and alternative pathway catecholamines showed that noradrenaline and adrenaline were more potent agonists than were octopamine, synephrine or deoxyadrenaline ($P < 0.05$). The stimulatory effects of noradrenaline and adrenaline (10^{-6} M) and of octopamine, synephrine and deoxyadrenaline (10^{-4} M) were inhibited by propranolol (10^{-5} M, $P < 0.05$, Table 1) but not by phentolamine (10^{-5} M, $P > 0.05$). Neither propranolol nor phentolamine significantly altered basal progesterone production (data not shown).

Discussion

These results demonstrate a role for the alternative pathway catecholamines octopamine, synephrine and deoxyadrenaline in the stimulation of bovine luteal steroidogenesis. The stimulatory effects of noradrenaline and adrenaline on progesterone production and the ability of the β -adrenergic antagonist propranolol to inhibit this response is consistent with previous results (Condon & Black, 1976; Jordan *et al.*, 1978; Harwood *et al.*, 1980; Ratner *et al.*, 1980; Norjavaara *et al.*, 1982). In other mammalian tissues octopamine has been reported to interact with α and β -adrenergic receptors (Fujiwara-Keisuke, Hideo, Takashi & Yoshiko, 1968), exclusively with α -receptors (Kelly & Burks, 1974; Kleinrok, 1979) or exclusively with β -adrenergic receptors (Chiba, 1976). The ability of propranolol but not phentolamine to inhibit the steroidogenic response to octopamine, synephrine and deoxyadrenaline suggests that their action is mediated by β -adrenergic receptors. The lower potency observed for the alternative pathway catecholamines with respect to progesterone production is similar to that reported for their ability to influence other physiological processes (Lands & Grant, 1952; Fujiwara *et al.*, 1968; Nedergaard & Westermann, 1968; Kelly & Burks, 1974).

The mammalian ovary is extensively innervated with adrenergic and cholinergic nerve fibres, although the CL appears to be devoid of any direct neural innervation (Bahr, Kao & Niswender, 1974; Burden, 1978; Stefenson, Owman, Sjöberg, Spörng & Wallies, 1981). No data are presently available to suggest that the bovine CL can synthesize or retain catecholamines. HPLC analysis of bovine luteal tissue did not

reveal noradrenaline or adrenaline to be present in early, mid or late cycle CL (P. Battista, C. Rexroad & W.A. Condon, unpublished observations). The failure to identify these catecholamines in bovine luteal tissue may suggest that the steroidogenic action of octopamine, synephrine and deoxyadrenaline is not due to conversion to primary catecholamines. Additionally, the inability of tyramine to stimulate luteal steroidogenesis may suggest that the enzymes responsible for primary and alternative pathway catecholamine synthesis are not present in bovine luteal tissue.

The mechanism whereby catecholamines stimulate progesterone production in the bovine is unknown. In rat granulosa cells, adrenergic agents increase the enzymic activity of 3 β -hydroxysteroid dehydrogenase while decreasing 20 α -hydroxysteroid dehydrogenase activity (Hsueh, *et al.* 1983). Ovarian denervation or chemical sympathectomy decreases 3 β -hydroxysteroid dehydrogenase activity in the rat CL, suggesting that catecholamines may regulate steroidal enzyme activity *in vivo* (Burden, 1978).

In summary, these results demonstrate that the alternative pathway catecholamines octopamine, synephrine and deoxyadrenaline stimulate *in vitro* steroid biosynthesis in bovine luteal tissue. The response to these catecholamines is less than that observed for noradrenaline and adrenaline, and the steroidogenic response of octopamine, synephrine and deoxyadrenaline is mediated by β -adrenergic receptors.

Scientific Contribution No. 1406 from the New Hampshire Agricultural Experiment Station. We thank Dr G. D. Niswender for the progesterone antiserum and D. Stranger for manuscript preparation. This research was a contribution to Northeast Regional Project NE-72.

References

- Axelrod, J. (1963) Enzymatic formation of adrenaline and other catechols from monophenols. Science, N.Y. 140, 499-500.
- Axelrod, J. & Saavedra, J.M. (1977) Octopamine. Nature, Lond. 265, 501-504.
- Bahr, J., Kao, L. & Naibandov, A.V. (1974) The role of catecholamines and nerves in ovulation. Biol. Reprod. 10, 273-290.
- Baldessarini, R.J. (1971) Release of aromatic amines from brain tissues of the rat in vitro. J. Neurochem. 18, 2509-2518.
- Baldessarini, R.J. & Vogt, M. (1971) The uptake and subcellular distribution of aromatic amines in the brain of the rat. J. Neurochem. 18, 2519-2533.
- Baldessarini, R.J. & Vogt, M. (1972) Regional release of aromatic amines from tissues of the rat brain in vitro. J. Neurochem. 19, 755-761.
- Battista, P.J. & Condon, W.A. (1986) Serotonin-induced stimulation of progesterone production by cow luteal cells in vitro. J. Reprod. Fert. 76, 231-238.
- Birnbaumer, L., Yang, P.C., Hunzicker-Dunn, M., Brockaert, J. & Duran, J.M. (1976) Adenyl cyclase activities in ovarian tissues. I. Homogenization and conditions of assay in Graafian follicles and corpora lutea of rabbits, rats and pigs: regulation by ATP, and some comparative properties. Endocrinology 99, 163-184.
- Burden, H.W. (1978) Ovarian innervation. In The Vertebrate Ovary, pp. 615-638. Ed. R. E. Jones. Plenum Press, New York.
- Chiba, S. (1976) Pharmacologic analysis of positive chronotropic and inotropic responses to octopamine. Tohoku J. exp. Med. 118, 247-253.
- Cooper, J.R., Bloom, F.E. & Roth, R.H. (1982) Catecholamines I: General aspects. In The Biochemical Basis of Neuropharmacology, pp. 109-172. Oxford University Press, New York.
- Condon, W.A. & Black, D.L. (1976) Catecholamine-induced stimulation of progesterone by the bovine corpus luteum in vitro. Biol. Reprod. 15, 573-578.
- Fischer, J.E., Musacchio, J., Kopin, I.J. & Axelrod, J. (1964) Effects of denervation on the uptake and β -hydroxylation of tyramine in the rat salivary gland. Life Sci. 3, 413-419.
- Fischer, J.E., Horst, W.D. & Kopin, I.J. (1965) β -Hydroxylated sympathomimic amines as false neurotransmitters. Br. J. Pharmacol. Chemother. 24, 477-484.

- Fujiwara, M., Keisuke, H., Hideo, M., Takashi, M. & Yoshiko, K. (1968) Pharmacological action of octopamine with special reference to biochemical conversion to noradrenaline. Jap. J. Pharmacol. 18, 113-129.
- Godkin, J.D., Black, D.L. & Duby, R.T. (1977) Stimulation of cyclic AMP and progesterone synthesis by LH, PGE₂ and isoproterenol in the bovine CL in vitro. Biol. Reprod. 17, 514-518.
- Harmar, A.J. & Horn, A.S. (1976) Octopamine in mammalian brain: Rapid post mortem increase and effects of drugs. J. Neurochem. 26, 987-993.
- Harwood, J.P., Dufau, M.L. & Catt, K.J. (1979) Differing specificities of ovarian cyclase by epinephrine and human chorionic gonadotropin. Molec. Pharmacol. 15, 439-444.
- Harwood, J.P., Richert, N.D., Dufau, M.L. & Catt, K.J. (1980) Gonadotropin-induced desensitization of epinephrine action in the luteinized rat ovary. Endocrinology 107, 280-288.
- Hsueh, A.J.W., Jones, P.B.C., Adashi, E.Y., Wang, C., Zhuang, L.Z. & Welsh, T.H., Jr (1983) Intraovarian mechanisms in the hormonal control of granulosa cell differentiation in rats. J. Reprod. Fert. 69, 325-342.
- Jordan, A.W., III, Caffrey, J.L. & Niswender, G.D. (1978) Catecholamine-induced stimulation of progesterone and adenosine 3',5'-monophosphate production by dispersed ovine luteal cells. Endocrinology 103, 385-392.
- Kakimoto, Y. & Armstrong, M.D. (1962) On the identification of octopamine in mammals. J. Biol. Chem. 237, 422-427.
- Kelly, R.J. & Burks, T.F. (1974) Relative vasoconstrictive potencies of norepinephrine, alpha-methyl-norepinephrine and octopamine. Arch. Int. Pharmacodyn. Ther. 208, 306-316.
- Kleinrok, Z. (1979) Central effects of octopamine administered into the lateral ventricle of rats. Acta physiol. pol. 30, 445-454.
- Kopin, I.J., Fischer, J.E., Musacchio, J. & Horst, W.D. (1964) Evidence for a false neurochemical transmitter as a mechanism for the hypotensive effect of monoamine oxidase inhibitors Proc. natn. Acad. Sci. U.S.A. 52, 716-721.
- Kopin, I.J., Fischer, J.E., Musacchio, J., Horst, W.D. & Weise, V.K. (1965) False neurochemical transmitters and the mechanism of sympathetic blockage by monoamine oxidase inhibitors. J. Pharmacol. exp. Therap. 147, 186-193.
- Lands, A.M. & Grant, J.I. (1952) The vasopressor action and toxicity of cyclohexylethylamine derivatives. J. Pharmacol. exp. Ther. 106, 341-345.

- Milvae, R.A., Allia, H.W. & Hansel, W. (1983) Methylation in bovine luteal cells as a regulator of luteinizing hormone action. Biol. Reprod. 29, 849-855.
- Mollinoff, P. & Axelrod, J. (1969) Octopamine: normal occurrence in sympathetic nerves of rats. Science, N.Y. 164. 428-429.
- Mollinoff, P.B. & Axelrod J. (1972) Distribution and turnover of octopamine in tissues. J. Neurochem. 19, 157-163.
- Musacchio, J., Kopin, I.J. & Snyder, S. (1964) Effects of disulfiram on tissue norepinephrine content and subcellular distribution of dopamine, tyramine and their β -hydroxylated metabolites. Life Sci. 3, 769-775.
- Nedergaard, O.A. & Westermann, E. (1968) Action of various sympathetic amines on the isolated stripped vas deferens of the guinea pig. Br. J. Pharmacol. 34, 475-483.
- Norjavaara, E., Seistam, G. & Ahren, K. (1982) Catecholamine stimulation of cyclic AMP and progesterone production in rat corpora lutea of different ages. Acta endocr., Copenh. 100, 613-622.
- Pate, J.L & Condon, W.A. (1982) Effect of serum and lipoproteins on steroidogenesis in cultured bovine luteal cells. Molec. cell. Endocr. 28, 551-562.
- Ratner, A., Sanborn, C.R. & Weiss, G.K. (1980) β -adrenergic stimulation of cAMP and progesterone in rat ovarian tissue. Am. J. Physiol. 239, E139-E143.
- Saavedra, J.M. (1974) Enzymatic-isotope method for octopamine at the picogram level. Analyt. Biochem. 59. 628-633.
- Snyder, S.H., Michaelson, I.A. & Musacchio, J. (1964) Purification of norepinephrine storage granules from rat heart. Life Sci. 3, 965-970.
- Snyder, S.H., Glowinski, J. & Axelrod, J. (1965) The storage of norepinephrine and some of its derivatives in brain synaptosomes. Life Sci. 4, 797-807.
- Stefenson, A., Owman, Ch., Sjöberg, N.O., Spörng, B. & Wallis, B. (1981) Comparative study of the autonomic innervation of the mammalian ovary, with particular regard to the follicular system. Cell Tissue Res. 215, 47-62.

Table 1. Effect of α - and β -adrenergic antagonists on primary and alternative pathway catecholamine-induced progesterone production (ng/ml incubation medium)

	N	Octopamine	Synephrine	Deoxyadrenaline	Noradrenaline	Adrenaline
Agonist alone	6	90.6 \pm 2.25 ^a	87.4 \pm 4.81 ^a	96.5 \pm 1.92 ^a	99.9 \pm 6.61 ^a	107.2 \pm 4.07 ^a
Agonist + propranolol	4	57.8 \pm 8.54 ^b	61.4 \pm 5.77 ^b	52.3 \pm 6.18 ^b	55.5 \pm 3.30 ^b	63.8 \pm 6.64 ^b
Agonist + phentolamine	4	93.9 \pm 4.87 ^a	87.9 \pm 7.88 ^a	89.8 \pm 5.42 ^a	89.3 \pm 6.18 ^a	98.7 \pm 4.34 ^a


Values are mean \pm s.e.m.


Means within columns with different superscript letters are significantly different ($P < 0.05$).


Control value equals 46.4 \pm 2.13.

Figure Legends

- Fig. 1. Primary and alternative pathways for the formation of catecholamines: (1) tyrosine hydroxylase; (2) aromatic aminoacid decarboxylase; (3) dopamine- β -oxidase; (4) phenylethylamine-N-methyl transferase; (5) nonspecific N-methyltransferase; (6) catechol-forming enzyme. Solid lines denote primary pathway. (Taken from Axelrod, 1963; and Cooper, Bloom & Roth, 1982).
- Fig. 2. Effect of alternative pathway catecholamines on progesterone production by dissociated bovine luteal cells. Data are expressed as the mean \pm s.e.m. (N=6 for all treatments). Solid line denotes control value of 46.4 ± 2.13 ng/ml. Bars within treatments with different superscript letters are significantly different ($P < 0.05$).
- Fig. 3. Effect of noradrenaline and adrenaline on progesterone production by dissociated luteal cells. Data are expressed as the mean \pm s.e.m. (N=6 for all treatments). Solid line denotes control value of 46.4 ± 2.13 ng/ml. Bars within treatments with different superscript letters are significantly different ($P < 0.05$).







EXPERIMENT FOUR

Biogenic amine stimulation of bovine luteal
progesterone production in vivo

P.J. Battista, J.P. Poff, D.R. Deaver¹ and W.A. Condon

Department of Animal and Nutritional Sciences
The University of New Hampshire
Durham, New Hampshire 03824, U.S.A.

and

Dairy Breeding Research Center¹
Pennsylvania State University
University Park, Pennsylvania 16802, U.S.A.

Accepted for publication in the Journal
of Reproduction and Fertility

Summary

The purpose of the present study was to determine if the administration of noradrenaline, adrenaline and serotonin could stimulate luteal progesterone production in vivo, as has been observed in vitro. Biogenic amines were administered using osmotic pumps placed subcutaneous in the neck region of regularly cycling, nonlactating dairy cows on days 9-11 (oestrus = Day 0) of the oestrous cycle. To monitor the effects of biogenic amines on luteal function, blood samples were collected using indwelling jugular catheters and the plasma fraction analyzed for progesterone. Samples were collected at 4h intervals for the first 12h of treatment and thereafter at 12h intervals for the remainder of the 72h treatment period. Upon administration of varying doses of noradrenaline, adrenaline and serotonin (0.5-2.0ug/kg/h) significant elevation of plasma progesterone concentrations was achieved at a dosage of 2.0 ug/kg/h ($P < 0.01$). The response to adrenaline was greater than that observed for noradrenaline and serotonin ($P < 0.05$). Within treatment comparison to pretreatment samples showed plasma progesterone concentrations to increase within 4h after the administration of noradrenaline, adrenaline and serotonin ($P < 0.05$) and this enhancement was maintained throughout the treatment period ($P < 0.05$). The elevation in plasma progesterone concentrations induced by noradrenaline, adrenaline and serotonin was independent of changes in circulating levels of luteinizing hormone. These results support a physiological role for endogenous biogenic amines in the enhancement of bovine luteal progesterone production.

Introduction

The catecholamines, noradrenaline and adrenaline, can stimulate the in vitro biosynthesis of cAMP and progesterone by luteal tissue from a variety of species (cow: Condon & Black, 1976; Godkin et al., 1977; Milvae, et al. 1983; rat: Harwood et al., 1979; Ratner et al., 1980; Norjavaara et al., 1982; rabbit: Birnbaumer et al., 1976; ewe: Jordan et al., 1978; and sow: Perkins et al., 1986). Recently we reported that the indoleamine serotonin enhances the production of progesterone by bovine luteal cells in vitro (Battista & Condon, 1986).

In vivo, the infusion of noradrenaline, adrenaline or the β -adrenergic agonist isoproterenol elevated plasma progesterone concentrations in the ewe (Bolt & Rollins, 1976), and infusion of noradrenaline increased cAMP levels in corpora lutea of the rat (Norjavaara et al., 1983). In contrast, in vivo administration of noradrenaline and isoproterenol was ineffective in stimulating progesterone production in oestrous rats, although fenoterol, a β_2 -adrenergic agonist, resulted in enhanced progesterone production (Zsolnai et al., 1982). Infusion of adrenaline in early pregnant women increased plasma progesterone (Fylling, 1971a), but these effects may be due to stimulation of placental progesterone production rather than a direct effect on luteal function (Fylling, 1971b; Flint et al., 1974; Csapo & Herczeg, 1977). This is supported by the observation that human luteal tissue is refractory to stimulation by exogenous adrenergic agonists in vitro (Richardson & Masson, 1980; Casper & Cotterell, 1984).

The infusion of isoproterenol or the β -adrenergic antagonist propranolol in pseudopregnant rabbits did not alter luteal function

(Gadsby et al., 1985). These authors suggested that endogenous catecholamines are not involved in the in vivo stimulation of luteal steroidogenesis. The physiological significance, if any, of biogenic amines in the in vivo enhancement of bovine luteal progesterone production is unknown. It was the purpose of this study to determine if the administration of noradrenaline, adrenaline, and serotonin could stimulate the production of progesterone in vivo, as has been observed in vitro.

Materials and Methods

Experimental Procedures

Serotonin, L-noradrenaline and L-adrenaline (Sigma Chemical Co., St. Louis, MO) were administered using osmotic pumps (Alza Corp., Palo Alto, CA). Pumps were inserted subcutaneously in the neck region of regularly cycling, nonlactating dairy cows on day 9-11 of the oestrous cycle (oestrous = Day 0) under general (Rompun, Miles Laboratories, Inc., Shawnee, KA) and local (Lidocaine HCl, J.A. Webster, Inc., N. Billerica, MA) anaesthesia. Treatments were prepared in a 0.15M NaCl solution (pH 4.0) to deliver controlled release rates of 0.5, 1.0 and 2.0 ug/kg/h. Control animals received osmotic pumps containing NaCl solution only. Blood samples were collected using indwelling jugular catheters (Abbocath- T. 14-gauge x 5 1/2", Abbot Hospitals, Inc., N. Chicago, IL) and samples were collected at 4h intervals for the first 12h of treatment and thereafter at 12h intervals for the remainder of the 72h treatment period. Duplicate pretreatment samples were obtained from each animal by jugular venipuncture prior to experimental manipulation. Blood samples were transferred to heparinized tubes and placed on ice for transport to the laboratory. Samples were centrifuged (4°C) for 20 min at 1.500 x g after which the plasma fraction was removed and stored at -20°C until analyzed for progesterone and luteinizing hormone.

Progesterone present in plasma samples was quantitated after extraction by radioimmunoassay as previously described (Battista et al., 1984). The progesterone antiserum (No. 337, Niswender) was prepared in sheep against 11 α -hydroxyprogesterone hemisuccinate

conjugated to bovine serum albumin. The progesterone tracer used was [1,2-³H] progesterone (New England Nuclear, Boston, MA). The sensitivity of the assay as determined by the lower 95% confidence limit of the maximum binding in the absence of any unlabelled progesterone was 0.07ng. The overall recovery of progesterone extracted from plasma was $92 \pm 1.5\%$. The intra- and interassay coefficients of variability were 5.1% and 9.4%, respectively. All samples from an individual animal were analyzed within the same assay and plasma progesterone concentrations are represented as the mean value of duplicate determinations not corrected for recovery. In all cases, n equals the number of individual animals tested.

To assess the effects of chronic administration of biogenic amines on the release of LH, plasma samples were analyzed using a modification of the procedure described by Niswender *et al.* (1969). Purified ovine LH (LER-1056-C2) was radioiodinated using IODO-GEN (1,3,4,6-tetrachloro-3,6-disphenyl-glycouril - Pierce Chemical Co., Rockville, IL). The primary antibody (antiovine LH, GDN-15) was diluted 1:40,000 in phosphate-buffered saline (0.14M NaCl, 0.01M NaPO₄, pH 7.4) containing 2% normal rabbit serum. Second antibody precipitation was achieved using sheep-anti-rabbit sera (DBRC-1) diluted 1:8 in phosphate-buffered saline. Precipitation of the second antibody complex was facilitated using 10% polyethylene glycol. Plasma samples were assayed in duplicate using 100ul of plasma. The mean of duplicate values was used for each unknown and all samples were analyzed within one assay. The sensitivity of the assay averages 0.2 ng. The minimal detectable value was used for samples with nondetectable LH concentration. The intra-assay coefficient of variation was 9.5%.

Statistical Analysis

Differences between treatment means were evaluated using one-way analysis of variance and Student-Newman-Keuls mean separation procedure. Within treatments, differences were determined using paired t-test.

Results

The effect of biogenic amines on mean plasma progesterone concentration over the 72h treatment period are shown in Table 1. The administration of noradrenaline, adrenaline and serotonin at 0.5 and 1.0 ug/kg/h did not significantly alter plasma progesterone concentrations from those observed in animals receiving no treatment ($P>0.05$). Administration of biogenic amines at 2.0 ug/kg/h significantly elevated circulating levels of progesterone above those in nontreated animals ($P<0.01$). The response to adrenaline was greater than that found for noradrenaline and serotonin ($P<0.05$). The time-course response to biogenic amine stimulation is shown in Fig. 1. Administration of noradrenaline and adrenaline elevated plasma progesterone within 4h ($P<0.05$, Fig. 1a), and this stimulation was maintained throughout the treatment period ($P<0.05$). Similarly treatment with serotonin significantly elevated progesterone within 4h ($P<0.05$, Fig. 1b), and this elevation was likewise maintained throughout the 72h treatment period ($P<0.05$).

None of the biogenic amines tested significantly altered circulating levels of LH. Mean plasma LH concentrations over the 72h treatment period for noradrenaline, adrenaline, serotonin and control animals were 0.22 ± 0.11 , 0.22 ± 0.12 , 0.22 ± 0.11 and 0.26 ± 0.13 ng/ml, respectively ($P>0.05$).

Discussion

These results demonstrate a role for noradrenaline, adrenaline and serotonin in the in vivo stimulation of bovine luteal progesterone production. None of the biogenic amines tested significantly altered circulating levels of LH, suggesting a direct effect of these amines on the corpus luteum. These results are consistent with the finding that none of the biogenic amines used cross the blood-brain barrier (Axelrod et al., 1959; Douglas, 1970) or significantly alter pituitary LH release under acute conditions (Kamberl & McCann, 1969; Schneider & McCann, 1969; Kamberl et al., 1970; Blake, 1976). These results support the in vitro stimulatory effects of noradrenaline, adrenaline (Condon & Black, 1976; Godkin et al., 1977; Milvae et al., 1983) and serotonin (Battista & Condon, 1986a) on bovine luteal progesterone production. The in vivo enhancement of luteal function by noradrenaline and adrenaline reported in the present study supports previous results obtained in the ewe (Boit & Rollins, 1976) and rat (Norjavaara et al., 1983).

Although the mammalian ovary has both adrenergic and cholinergic nerve fibres, the corpus luteum does not appear to be directly innervated by either neural system (Bahr et al., 1974; Burden, 1978; Stefenson et al., 1981). Support for a physiological role of the nervous system in the control of ovarian steroidogenesis is suggested by the finding that electrical stimulation of specific brain regions increased the concentration of estradiol and progesterone in ovarian venous blood of hypophysectomized-adrenalectomized, pro-oestrous rats (Kawakami et al., 1981). Sectioning of the superior ovarian nerve decreased estradiol and progesterone concentrations in ovarian venous

blood of pro-oestrous rats (Aguado & Ojeda, 1984) while electrical stimulation of the superior ovarian nerve enhanced luteal progesterone production in dioestrous rats through stimulation of β -adrenergic receptors (Weiss *et al.*, 1982).

In addition to the absence of direct adrenergic innervation, we have been unable to detect noradrenaline or adrenaline in bovine luteal tissue using HPLC analysis (Battista & Condon, 1986b). In contrast, we have found serotonin in bovine luteal tissue (Battista & Condon, 1986a). Possible sources of luteal serotonin include mast cells, blood platelets, storage of peripherally circulating serotonin or de novo synthesis. Histological determination of mast cells using Giemsa, toluidene blue and methylene blue staining techniques has failed to demonstrate the presence of mast cells in either early or mid-cycle luteal tissue (unpublished observations). Serotonin present in luteal tissue may be derived from peripherally circulating serotonin similar to that reported for rat testicular (Ellis *et al.*, 1972) and adrenal (Verhofstad & Jonsson, 1983) tissues. We are presently conducting immunohistochemical experiments to examine these possibilities and to determine the cellular location of serotonin. The mechanism whereby biogenic amines enhance bovine luteal progesterone production is unknown. *In vitro* studies using rat granulosa cells showed primary catecholamines increased the enzymatic activity of 3β -hydroxysteroid dehydrogenase while decreasing 20α -hydroxysteroid dehydrogenase activity (Hsueh *et al.*, 1983). Ovarian denervation or chemical sympathectomy decreases 3β -hydroxysteroid dehydrogenase activity in both the interstitial gland and corpus luteum of pregnant rats (Burden & Lawrence, 1977).

The ability of the primary pathway catecholamines noradrenaline and adrenalineto stimulate luteal progesterone production both in vivo and in vitro suggests a physiological role for endogenous adrenergic control of bovine luteal function. Additionally, the ability of serotonin to enhance the in vivo and in vitro production of progesterone and the presence of serotonin within bovine luteal tissue suggests a physiological role for serotonin as an intraovarian stimulator of luteal function.

Scientific Contribution No. 1456 from the New Hampshire Agricultural Experiment Station. We thank Dr. G.D. Niswender for the progesterone antiserum and D. Stranger and M. Coburn for manuscript preparation. This research was a contribution to Northeast Regional Project NE-72.

References

- Aguado, L.I. & Ojeda, S.R. (1984) Ovarian adrenergic nerves play a role in maintaining preovulatory steroid secretion. Endocrinology 114, 1944-1946.
- Axelrod, J., Weil-Malherbe, H. & Tomchick, R. (1959) The physiological disposition of ^3H -epinephrine and its metabolite metanephrine. J. Pharmacol. exp. Ther. 137, 251-256.
- Bahr, J., Kao, L. & Naibandov, A.V. (1974) The role of catecholamines and nerves in ovulation. Biol. Reprod. 10, 273-290.
- Battista, P.J., Rexroad, C.E. Jr & Williams, W.F. (1984) Effects of progesterone administration to dairy helpers on sensitivity of corpora lutea to $\text{PGF}_2\alpha$ and on plasma LH concentration. Theriogenology 22, 47-58.
- Battista, P.J., & Condon, W.A. (1986a) Serotonin-induced stimulation of progesterone production by cow luteal cells *in vitro*. J. Reprod. Fert. 76, 231-238.
- Battista, P.J. & Condon, W.A. (1986b) A role for alternative pathway catecholamines in the regulation of steroidogenesis in cow luteal cells. J. Reprod. Fert. 78, 275-280.
- Birnbaumer, L., Yang, P.C., Hunzicker-Dunn, M., Brockaert, J. & Duran, J.M. (1976) Adenyl cyclase activities in ovarian tissues. I. Homogenization and conditions of assay in Graffian follicles and corpora lutea of rabbits, rats and pigs: regulation by ATP, and some comparative properties. Endocrinology 99, 163-184.
- Blake, C.A. (1976) Effects of intravenous infusion of catecholamines on rat plasma luteinizing hormone and prolactin concentrations. Endocrinology 98, 99-104.
- Bolt, D.J. & Rollins, R. (1976) Increased plasma progesterone during infusion of epinephrine, norepinephrine or isoproterenol in luteal phase ewes. Biol. Reprod. Suppl. 1, abstr. 43.
- Burden, H.W. & Lawrence, I.E. (1977) The effects of denervation on the localization of $\Delta^5-3\beta$ -hydroxysteroid dehydrogenase activity in the rat ovary during pregnancy. Acta anat. 97, 286-290.
- Burden, H.W. (1978) Ovarian innervation. In The Vertebrate Ovary, pp. 615-638. Ed. R.E. Jones. Plenum Press, New York. Casper, R.F. & Cotterell, M.A. (1984) The effect of adrenergic and cholinergic agents on progesterone production by human corpus luteum *in vitro*. Am. J. Obstet. Gynecol. 148, 663-669.
- Condon, W.A. & Black, D.L. (1976) Catecholamine-induced stimulation of progesterone by the bovine corpus luteum *in vitro*. Biol. Reprod.

15, 573-578.

- Csapo, A.I. & Herczeg, J. (1977) Arrest of premature labor by Isosuprine. Am. J. Obstet. Gynecol. 129, 482-491.
- Douglas, W.W. (1970) Histamine and antihistamine; 5-hydroxytryptamine and antagonists. In The Pharmacological Basis of Therapeutics. 4th edn, pp. 645-662. Eds. L.S. Goodman & A. Gilman. Macmillan Co., New York.
- Ellis, L.C., Jaussl. A.W., Baptista, M.H. & Urry, R.L. (1972) Correlation of age changes in monoamine oxidase activity and androgen synthesis by rat testicular minced and teased-tubular preparation in vitro. Endocrinology 90, 1610-1618.
- Flint, A.P.F., Anderson, A.B.M. & Turnbull, A.C. (1974) Control of utero-ovarian and peripheral venous plasma progesterone by sympathomimic drugs in pregnant sheep. J. Endocr. 63, 253-254.
- Fylling, P. (1971a) Vasopressin-induced increase in peripheral plasma progesterone levels during early human pregnancy. Acta endocr., Copenh. 66, 273-282.
- Fylling, P. (1971b) Propranolol-blockade of vasopressin induced increase in plasma progesterone in early human pregnancy. Acta endocr., Copenh. 66, 283-288.
- Gadsby, J.E., Keyes, P.L., Schwartz, T.S., Bill, C.H., II & Lucchesi, B. (1985) Do catecholamines play a physiological role in regulating corpus luteum function in the pseudopregnant rabbit? Biol. Reprod. 32, 907-915.
- Godkin, J.D., Black, D.L. & Duby, R.T. (1977) Stimulation of cyclic AMP and progesterone synthesis by LH, PGE₂ and Isoproterenol in the bovine CL in vitro. Biol. Reprod. 17, 514-518.
- Harwood, J.P., Richert, N.D., Dufau, M.L. & Catt, J.C. (1980) Gonadotropin-induced desensitization of epinephrine action in luteinized rat ovary. Endocrinology 107, 280-288.
- Hsueh, A.J.W., Jones, P.B.C., Adashi, E.Y., Wang, C., Zhuang, L.Z. & Welsh, T.H., Jr. (1983) Intraovarian mechanisms in the hormonal control of granulosa cell differentiation in rats. J. Reprod. Fert. 69, 325-342.
- Jordan, A.W., III, Caffrey, J.L. & Niswender, G.D. (1978) Catecholamine-induced stimulation of progesterone and adenosine 3',5'-monophosphate production by dispersed ovine luteal cells. Endocrinology 103, 385-392.
- Kamberl, I.A. & McCann, S.M. (1969) Effect of biogenic amines, FSH-releasing factors (FRF) and other substances on the release of FSH by pituitaries incubated in vitro. Endocrinology 85, 815-824.
- Kamberl, I.A., Mical, R.S. & Porter, J.C. (1970) Effects of anterior

- pituitary perfusion and intraventricular injection of catecholamines on LH release. Endocrinology 87, 1-12.
- Kawakami, M., Kubo, K., Uemura, T., Nagase, M. & Hayashi, R. (1981) Involvement of ovarian innervation in steroid secretion. Endocrinology 109, 136-145.
- Milvae, R.A., Ailla, H.W. & Hansel, W. (1983) Methylation in bovine luteal cells as a regulator of luteinizing hormone action. Biol. Reprod. 29, 849-855.
- Niswender, G.D., Reichert, L.E., Jr., Midgley, A.R., Jr. & Nalbandov, A.V. (1969) Radioimmunoassay for bovine and ovine luteinizing hormone. Endocrinology 84, 1116-1173.
- Norjavaara, E., Selstam, G. & Ahren, K. (1982) Catecholamine stimulation of cyclic AMP and progesterone production in rat corpora lutea of different ages. Acta endocr. Copenh. 100, 613-622.
- Norjavaara, E., Selstam, G., Damberg, J.E. & Johansson, B.M. (1983) In vivo effect of noradrenaline on the cyclic AMP level in rat corpora lutea. Acta physiol., Scand. 119, 113-116.
- Perkins, S.N., Cronin, M.J. & Veldhuis, J.D. (1986) Properties of β -adrenergic receptors on porcine corpora lutea and granulosa cells. Endocrinology 118, 998-1005.
- Ratner, A., Sanborn, C.R. & Weiss, G.K. (1980) β -Adrenergic stimulation of cAMP and progesterone in rat ovarian tissue. Am. J. Physiol. 239, E139-E143.
- Richardson, M.C. & Masson, G.M. (1980) Progesterone production by dispersed cells from human corpus luteum: stimulation by gonadotrophins and prostaglandin $F_{2\alpha}$. Lack of response to adrenaline and isoprenaline. J. Endocr. 87, 247-254.
- Schneider, H.P.G. & McCann, S.M. (1969) Possible role of dopamine as transmitter to promote discharge of LH-releasing factor. Endocrinology 85, 121-132.
- Stefenson, A., Owman, Ch., Sjöberg, N.O., Spörng, B. & Waller, B. (1981) Comparative study of the autonomic innervation of the mammalian ovary, with particular regard to the follicular system. Cell Tissue Res. 215, 47-62.
- Verhofstad, A.A.J. & Jonsson, G. (1983) Immunohistochemical and neurochemical evidence for the presence of serotonin in the adrenal medulla of the rat. Neuroscience 10, 1443-1453.
- Weiss, G.K., Dall, W.G. & Ratner, A. (1982) Evidence for direct neural control of ovarian steroidogenesis in rats. J. Reprod. Fert. 65, 507-511.
- Zsolnai, B., Varga, B. & Horvath, E. (1982) Increase of ovarian progesterone secretion by β_2 -adrenergic stimulation in oestrous

rats. Acta endocr. Copenh. 101, 268-272.


TABLE 1: Effect of biogenic amines on mean plasma progesterone concentrations (ng/ml) over a 72h treatment period.


Biogenic Amine	N	Concentration (ug/kg/h)	Plasma Progesterone (mean \pm s.e.m.)
Noradrenaline	2	0.5	4.0 \pm .37 ^a
	2	1.0	4.8 \pm .40 ^a
	4	2.0	7.9 \pm .26 ^b
Adrenaline	2	0.5	4.7 \pm .60 ^a
	2	1.0	5.1 \pm .62 ^a
	4	2.0	9.0 \pm .46 ^c
Serotonin	2	0.5	4.5 \pm .17 ^a
	2	1.0	4.4 \pm .29 ^a
	5	2.0	7.9 \pm .23 ^b
None	4	0	4.2 \pm .17 ^a

N equals the number of individual animals tested. Means with different superscript letters are significantly different ($P < 0.01$ for noradrenaline, adrenaline and serotonin vs. control, $P < 0.05$ for adrenaline vs. noradrenaline and serotonin).

FIGURE LEGEND

FIGURE 1. Time-course effect of (a) noradrenaline, adrenaline and (b) serotonin on the in vivo stimulation of bovine luteal progesterone production. Values are the mean + s.e.m., N = 4 for noradrenaline, adrenaline and control, N = 5 for serotonin. All treatment values are significantly different from corresponding time 0 values ($P < 0.05$).





SUMMARY AND SIGNIFICANCE OF RESULTS

Catecholamine Studies

- 1) The enhancement of bovine luteal P_4 production in response to catecholamines is mediated by a mixed population of β -adrenergic receptors. Both β_1 - and β_2 -adrenergic receptors are present on bovine luteal cell membranes.
- 2) The catecholamines: octopamine, synephrine and dexoyepinephrine, which are derived through an alternative or secondary pathway, stimulated basal P_4 production. The stimulatory effect of these amines was mediated by β -adrenergic receptors.
- 3) HPLC analysis of bovine luteal tissue did not demonstrate detectable levels of either NE or E in luteal tissue obtained at various stages of the estrous cycle.
- 4) The in vivo administration of NE and E stimulated bovine luteal function, as assessed by elevated levels of plasma P_4 . The increase in circulating levels of P_4 in response to NE and E treatment was independent of changes in circulating levels of LH.

Biogenic Amine Studies

- 1) The addition of 5-HT resulted in a dose-dependent enhancement of P_4

production by dissociated bovine luteal cells in short-term incubation.

2) Incubation of luteal cells with the 5-HT metabolite 5-MTA, also stimulated the production of P_4 .

3) The response to 5-HT was mediated by a 5-HT₁ receptor which may not be associated with adenylate cyclase.

4) Analysis of bovine luteal tissue by HPLC demonstrated the presence of 5-HT and showed that the concentration of 5-HT varied with the stage of luteal development.

5) Administration of 5-HT in vivo resulted in elevated levels of plasma P_4 . The change in plasma P_4 was independent of alterations in circulating levels of LH.

The ability of catecholamines to stimulate luteal steroidogenesis both in vitro and in vivo supports a physiological role for endogenous catecholamines in the stimulation of bovine luteal function. The inability to detect NE and E with bovine luteal tissue suggests that the bovine CL is unable to synthesize or store catecholamines. Thus, neurotransmitter input to the CL may arise from the sympathetic neural network present within the ovary or from circulating catecholamines. Direct experimentation of catecholamine synthesis by luteal tissue has not been conducted. Further research is required to address this question, and to determine the significance of innervation to the normal functioning of the bovine CL.

However, based on the results of the present studies it would be anticipated that inhibition of neural input to the CL would result in luteal insufficiency. Thus Intraovarian Infusion of a β -adrenergic antagonist such as propranolol, should result in reduced circulating levels of P_4 , thereby attributing physiological significance to neural stimulation in the normal functioning of the bovine CL. It could be speculated that a deficiency in neural input to the CL may in part account for pregnancy failure. Since successful conception is dependent upon adequate levels of P_4 , insufficient neural input might explain the lower levels of P_4 associated with some repeat-breeder dairy cattle.

Additionally, neural enhancement of bovine luteal function may be involved in explaining reproductive insufficiency resulting from stress. Stress factors such as extremes in temperature, milk production, nutritional deprivation, or excessive physical exercise may result in enhanced neural stimulation of luteal steroidogenesis. As a result, circulating levels of P_4 would be elevated above and beyond normal limits. Prolonged exposure to P_4 would then alter steroid feedback mechanisms at the hypothalamus and pituitary, thereby inhibiting reproductive cyclicity.

The ability of 5-HT to stimulate luteal P_4 production both in vitro and in vivo, coupled with the presence of 5-HT within bovine luteal tissue. Is supportive of a physiological role for endogenous 5-HT as an Intraovarian stimulator of luteal function. Further studies are required to directly access the serotonergic contribution to normal luteal function, to determine the cellular location of 5-HT within the CL, and to assess the capabilities for de novo synthesis.

With respect to physiological significance, the higher concentration

of 5-HT present in newly formed luteal tissue may suggest a role for 5-HT in the process of luteinization. 5-HT may also play a luteotropic role during early CL formation, a time at which LH is ineffective in stimulating P_4 production.

It would be anticipated based on the results of the present studies that the administration of a 5-HT antagonist such as mianserin, would reduce luteal P_4 secretion. Such an experiment would demonstrate more directly the serotonergic contribution to normal luteal function. If 5-HT antagonists exhibit luteolytic properties, such a pharmacological tool may lead to better methods for the induction of luteolysis, and thus better strategies for estrous cycle control.

Lastly, 5-HT may be involved in early pregnancy recognition. Based on the finding that embryos can secrete platelet activating factor and that platelets contain 5-HT. It is possible that embryo-derived platelet activating factor may act on platelets within the CL to stimulate the release of 5-HT, which would then stimulate P_4 secretion by the maternal CL. Thus, identification of the mechanisms controlling early pregnancy recognition may prove helpful in preventing early embryonic death.

In summary, the results of these studies support the concept that biogenic amines contribute to the physiological control of bovine luteal function. In the future these results may lead to new areas of chemical control of luteal function, which may result in better synchronization strategies for breeding and embryo transfer.